

Article

Roles of physical activity in improving both physiological and psychological health outcomes among thyroid cancer survivors

Yang Wang^{1,*}, Zainal Abidin B Zainuddin², Haitao Yin¹

¹ Department of Physical Education, Yuncheng University, 044000 Yuncheng, Shanxi, China

² Faculty of Educational Sciences and Technology, Universiti Teknologi Malaysia, 81310 Skudai, Johor, Malaysia

* Corresponding author: Yang Wang, wangyang@ycu.edu.cn

CITATION

Wang Y,Zai Z,Yin H.
Roles of physical activity in improving both physiological and psychological health outcomes among thyroid cancer survivors. *Psycho-Oncologie*. 2026; 20(3): 4174. <https://doi.org/10.18282/po4174>

ARTICLE INFO

Received: 28 October 2025

Accepted: 30 March 2026

Available online: Date Month Year

COPYRIGHT



Copyright © Year by author(s).
Psycho-Oncologie is published by PiscoMed Publishing Pte Ltd. This work is licensed under the Creative Commons Attribution (CC BY) license.
<https://creativecommons.org/licenses/by/4.0/>

Abstract: Background: Thyroid cancer is the most common endocrine malignancy, and survivors often face ongoing health challenges. Physical activity has been proposed as a non-pharmacological intervention with the potential to improve overall health and quality of life in cancer survivors, yet its role in thyroid cancer survivors is still under-explored. **Objective:** This systematic review aims to synthesize current evidence on the effects of physical activity on health outcomes in thyroid cancer survivors, focusing on both physiological and psychological aspects. **Methods:** A comprehensive search of PubMed, the Web of Science, and Scopus was conducted to identify studies published between 2013 and 2023. Studies were included if they focused on thyroid cancer survivors, incorporated physical activity interventions, and reported outcomes related to physical health, psychological well-being, or quality of life. **Results:** Physical activity was found to positively affect multiple health domains in thyroid cancer survivors. Physiological benefits included improved cardiorespiratory fitness, enhanced hormonal regulation (T3, T4, and TSH), and reduced fatigue. Psychological outcomes, such as reduced anxiety and depression, were consistently reported alongside improvements in overall quality of life. Evidence suggests that moderate-intensity aerobics and yoga may be particularly effective, although variations in intervention duration and intensity were observed across studies. **Conclusion:** Physical activity offers substantial physical and psychological benefits for thyroid cancer survivors. Future large, well-designed trials are needed to determine optimal exercise protocols and long-term effects. Tailored physical activity programs integrated into survivorship care can improve recovery and quality of life.

Keywords: thyroid cancer, physical activity, health outcomes, psychological well-being, systematic review

1. Introduction

1.1. Background and significance

Thyroid cancer has emerged as the most prevalent endocrine malignancy worldwide, with its incidence steadily rising over recent decades[1]. Advancements in diagnostic methods, such as high-resolution ultrasound and fine-needle aspiration biopsy, have enabled earlier detection and generally lead to favorable prognoses[2]. Despite high survival rates—especially in differentiated thyroid cancers like papillary and follicular subtypes—many survivors continue to face persistent challenges following treatment[3,4], including chronic fatigue, reduced cardiorespiratory fitness, altered metabolic status, and anxiety about disease recurrence, alongside the psychological strain of maintaining thyroid hormone suppression therapy[5,6].

Increasingly, physical activity is recognized as a potential intervention to address

these challenges in oncology care[7]. Evidence from various cancer populations shows that exercise can significantly improve physical functioning, mitigate treatment-related side effects, and enhance quality of life[8]. In breast and colorectal cancers, structured exercise programs have been linked to better cardiorespiratory endurance, less fatigue, and reduced depressive symptoms[9]. However, the specific impact of exercise on thyroid cancer survivors remains underexplored. This is particularly important given the distinct clinical characteristics of thyroid cancer survivors, including prolonged reliance on thyroid hormone replacement and altered metabolic function, which may affect their response to exercise interventions.

Moreover, the psychosocial aspects of thyroid cancer survivorship warrant increasing attention[10,11]. Survivors often experience heightened stress and uncertainty related to disease progression, secondary cancer risks, and the need for lifelong monitoring[12,13]. Physical activity's potential to mitigate these emotional burdens, including anxiety and depression, has been observed in other cancer contexts, but its relevance to thyroid cancer survivors requires further investigation[5,14]. Understanding how exercise can enhance both physical recovery and emotional well-being is essential for optimizing care and improving overall quality of life in this population.

1.2. Necessity of the review

While substantial evidence from general cancer survivorship research highlights the benefits of physical activity, applying these findings directly to thyroid cancer survivors is challenging due to the disease's distinct pathophysiology and treatment protocols[15,16]. Thyroid cancer patients often undergo thyroidectomy, resulting in lifelong dependency on thyroid hormone replacement[17]. Additional treatments, such as radioactive iodine therapy, may further impact metabolic rate, body composition, and energy levels in ways not commonly observed in other cancer types[18]. These factors suggest that exercise prescriptions for thyroid cancer survivors may need to be adjusted in terms of intensity, frequency, and type compared to recommendations for other cancer populations[19].

Moreover, the body of literature on physical activity among thyroid cancer survivors remains limited, often characterized by small sample sizes, varied study designs, and inconsistent outcome measures [20]. While some studies focus on biochemical markers like TSH, T3, and T4 levels, others examine quality of life, psychological well-being, or fatigue reduction [21,22]. This variability makes it difficult to establish a clear, unified understanding of how physical activity interventions benefit this group. A comprehensive review that synthesizes existing studies is essential not only to summarize current findings but also to highlight gaps in research methodologies and intervention strategies.

The growing emphasis on individualized survivorship care further underscores the need for tailored, evidence-based exercise recommendations[23]. Thyroid cancer survivors differ in age, comorbidities, and the extent of their disease and treatment[24,25]. Therefore, general exercise guidelines may not adequately address the diverse needs of this population[26]. By systematically reviewing the available evidence, we can better identify the most beneficial types of exercise—whether aerobic, resistance, or yoga—along with the appropriate intensities (low, moderate, or high) and durations that optimize both physical and psychological outcomes.

1.3. Research objectives

Given the increasing number of thyroid cancer survivors and the significant gap

in targeted research, this systematic review aims to achieve several key objectives:

Evidence Synthesis: To collate and summarize the existing studies on the impact of physical activity on health outcomes in thyroid cancer survivors. This includes a focus on both physiological markers (such as hormonal balance, metabolic function, and fatigue) and psychological well-being (including anxiety, depression, and quality of life)[27].

Comparison of Interventions: To evaluate the effectiveness, feasibility, and safety of various exercise interventions—ranging from aerobic training and resistance exercises to yoga and combined modalities—within this population[28].

Identification of Gaps and Future Directions: To pinpoint methodological limitations and gaps in the current literature, providing direction for future studies. We will highlight the need for larger randomized controlled trials, longer follow-up periods, and more refined outcome measures to help establish evidence-based exercise recommendations for thyroid cancer survivors.

Clinical and Policy Implications: By offering a comprehensive overview of how physical activity can influence recovery, this review seeks to inform clinical guidelines, survivorship programs, and health policy initiatives[29]. Empowering survivors with well-supported, evidence-based exercise interventions could significantly enhance their long-term health and well-being.

2. Materials and methods

2.1. Literature search strategy

A comprehensive literature search was conducted across three electronic databases—PubMed, Web of Science, and Scopus—covering publications from January 2013 to December 2023. The search combined Medical Subject Headings (MeSH) with relevant free-text keywords related to thyroid cancer, physical activity, exercise, health outcomes, and psychological health.

To ensure thoroughness, we also reviewed the reference lists of included studies to identify any additional relevant research. Only full-text, peer-reviewed articles were considered in the initial pool of studies. Conference abstracts, dissertations, and other grey literature were included on a case-by-case basis if they provided sufficient methodological details to evaluate quality and outcomes. No geographical restrictions were applied; however, due to resource and language constraints, only articles published in English were included.

2.2. Inclusion and exclusion criteria

All retrieved studies underwent a two-step screening process: first, a title/abstract review, followed by a full-text assessment. The eligibility criteria for inclusion and exclusion were as follows:

Inclusion Criteria: Studies involving thyroid cancer survivors (any histological subtype, including papillary, follicular, medullary, and anaplastic) who were either post-treatment or undergoing survivorship care. Physical activity or exercise as the primary intervention, or as a key component of a multi-modal program (e.g., aerobic exercise, resistance training, yoga, or other structured exercises). At least one physical or psychological health indicator, such as hormonal levels (TSH, T3, T4), fatigue scores, cardiorespiratory fitness, anxiety, depression, or quality of life measures[30]. Original research articles providing empirical data on the effects of the intervention.

Exclusion Criteria: Studies focusing on other cancer types or general oncology populations without specific subgroup analysis for thyroid cancer[31]. Studies that

did not include a defined physical activity or exercise intervention (e.g., purely dietary interventions or pharmacological treatments)[32]. Narrative reviews, systematic reviews, meta-analyses, editorials, letters, or commentaries lacking primary data[33]. Insufficient Outcome Data: Studies that did not report relevant physiological or psychological health outcomes.

Titles and abstracts were screened according to these criteria. Articles that met the inclusion criteria, or those with uncertain relevance, were retrieved for full-text review.

2.3. Data extraction and analysis

A standardized data extraction form was created to ensure consistent information collection from each included study. The following variables were documented:

General Study Information: Author(s), publication year, study design (e.g., randomized controlled trial, observational), and study location. Participant Characteristics: Sample size, mean or median age, sex distribution, thyroid cancer subtype, and treatment background (e.g., surgery type, use of radioactive iodine). Intervention Details: Type(s) of physical activity (aerobic exercise, resistance training, yoga), frequency (times per week), intensity (low, moderate, or high), duration (minutes per session and total intervention length), and any progression protocols specified[34]. Outcomes Measured: Physiological markers(e.g., T3, T4, TSH levels, cardiorespiratory fitness), psychological measures (e.g., anxiety, depression, quality of life scales), and secondary endpoints (e.g., body composition changes, fatigue indices)[35]. Main Findings: Statistical significance, effect sizes (if reported), and the authors' primary conclusions regarding the exercise intervention's impact.

Data were synthesized using a narrative approach, with a focus on the consistency of findings across studies. Subgroup analyses, where applicable—such as differences between exercise modalities or intensity levels—were noted. While the primary focus was on textual synthesis, concise tables were included to summarize key information (e.g., intervention types, outcome measures, and main results), facilitating cross-study comparisons and highlighting methodological patterns.

2.4. Quality assessment

All included studies were evaluated for methodological quality following the guidelines set by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA). Specifically:

Risk of Bias: Randomized controlled trials were assessed using established tools to evaluate domains such as sequence generation, allocation concealment, blinding, incomplete outcome data, and selective reporting. Observational studies were assessed using the Newcastle-Ottawa Scale (NOS)[36]. Study Heterogeneity: Variations in intervention duration, exercise intensity, and participant characteristics were documented as potential sources of heterogeneity[37]. Reporting Clarity: We examined each study's adherence to key reporting details, such as participant flow diagrams, baseline characteristics, and outcome definitions, to assess transparency and reproducibility.

Discussions and consensus meetings were held to resolve any disagreements in the quality assessment process, ensuring a robust and transparent evaluation. Studies with a higher risk of bias were not excluded but were critically appraised in the Results and Discussion sections, enabling readers to interpret the findings with appropriate caution.

3. Results and discussion

3.1. Literature search outcomes

A total of 1,513 records were initially retrieved from three databases (PubMed: 562; Web of Science: 480; Scopus: 471) using predefined search terms and filters (2013–2023). After removing 436 duplicate entries, 1,077 articles remained for title and abstract screening. Of these, 1,012 were excluded for not meeting the inclusion criteria (e.g., studies not focused on thyroid cancer or lacking an exercise component), leaving 65 articles for full-text review. Following the full-text assessment, an additional 50 articles were excluded for the following reasons:

Inadequate focus on exercise interventions (n = 21); Non-empirical articles (n = 13); Outcomes unrelated to physical or psychological health indicators (n = 9); Combined cancer populations without a specific subgroup analysis for thyroid cancer (n = 7).

Ultimately, 15 studies met the full inclusion criteria and were included in this systematic review. Figure 1 presents the PRISMA flow diagram, outlining the screening and selection process.

Characteristics of Included Studies

The 15 studies included in this review employed a range of research designs: 8 randomized controlled trials (RCTs), 4 prospective cohort or longitudinal studies, and 3 cross-sectional or quasi-experimental studies. Sample sizes varied from 28 to 248 participants, with a mean age ranging from the early 40s to late 50s. The majority of participants had undergone total or near-total thyroidectomy, often followed by radioactive iodine therapy, and most were receiving thyroid hormone replacement. Geographically, the studies spanned North America, Europe, and Asia, though the majority were conducted in Western countries.

A significant portion of the studies utilized structured exercise interventions, predominantly moderate-intensity aerobic or combined aerobic and resistance training regimens. Intervention durations ranged from 6 to 24 weeks, with session frequencies varying from 2 to 5 times per week. Several studies also explored alternative or complementary approaches, such as yoga or Pilates, assessing their feasibility and effects on both physical and psychological outcomes.

3.2. Summary of health-related outcomes

To facilitate clarity and depth of analysis, the results have been divided into three major outcome categories: (1) Physiological Health, (2) Psychological Health, and (3) Relationship Between Intervention Characteristics and Effects.

3.2.1. Physiological health

3.2.1.1. Thyroid hormone regulation and metabolic markers

Thyroid Hormones (T3, T4, TSH): Six of the fifteen studies included in this review assessed changes in thyroid hormone levels as primary or secondary endpoints[38]. Three randomized controlled trials reported slight improvements in the regulation of T3, T4, and TSH following moderate-intensity aerobic exercise performed three times per week over 8–12 weeks. While the observed changes were generally modest and did not always reach statistical significance, the authors suggested that consistent physical activity might contribute to better metabolic homeostasis, particularly when coupled with appropriate thyroid hormone replacement therapy[38].

Metabolic Parameters (Glucose, Lipid Profiles): Four studies examined

metabolic markers, including blood glucose and lipid profiles. The findings were mixed: two trials observed modest improvements in fasting glucose and cholesterol levels in the exercise groups compared to controls[39]. However, one quasi-experimental study found no significant changes in lipid profiles, attributing the lack of effect to the relatively short 6-week intervention duration[40].

3.2.1.2. Physical fitness and cardiorespiratory capacity

Cardiorespiratory Fitness: Ten studies evaluated cardiorespiratory fitness using VO2 max or other measures such as the 6-minute walk test. The majority found that structured aerobic exercise programs—such as brisk walking, cycling, or treadmill workouts—resulted in significant improvements in aerobic capacity[41]. For example, in one randomized controlled trial involving 120 thyroid cancer survivors, the intervention group (12 weeks of supervised brisk walking) showed a 15% increase in VO2 max, compared to a 4% increase in the control group ($p < 0.01$) [42].

Muscular Strength and Endurance: Five studies examined the effects of resistance training or combined aerobic-resistance regimens on muscular strength and endurance[43]. Overall, improvements in muscular endurance and strength were observed. However, three studies highlighted the importance of caution when prescribing resistance loads, especially for survivors with comorbid musculoskeletal conditions or post-surgical complications.

3.2.1.3. Exercise modality-specific physiological effects

Comparisons between different exercise modalities (e.g., aerobic vs. resistance, home-based vs. supervised) indicated that combined aerobic-resistance programs generally produced more comprehensive improvements in both physical function and body composition compared to aerobic exercise alone[44]. On the other hand, alternative approaches such as yoga and Pilates led to modest gains in flexibility and core strength, along with some reductions in stress levels, as discussed in more detail in Section 5.2.2[45].

3.2.2. Psychological health

3.2.2.1. Improvements in anxiety, depression, and mood

Anxiety and Depression: Nine studies utilized validated scales, including the Hospital Anxiety and Depression Scale (HADS) and the Beck Depression Inventory (BDI)[46]. These studies consistently showed that participants engaging in exercise interventions experienced significant reductions in anxiety and depression scores compared to control groups. For example, one randomized controlled trial ($n = 80$) reported a 25% reduction in HADS-Anxiety scores among the exercise group, compared to a 10% decrease in the usual-care group ($p < 0.05$)[47].

Stress and Mood Disturbances: Exercise programs incorporating mindfulness components, such as yoga, demonstrated particular effectiveness in reducing stress and mood disturbances. These interventions appeared to not only provide physical benefits but also enhance emotional regulation, potentially through mechanisms like focused breathing and the relaxation response[48].

3.2.2.2. Quality of life and daily functioning

Quality of Life (QoL) Assessments: The SF-36 (Short Form Health Survey) and the EORTC QLQ-C30 (European Organization for Research and Treatment of Cancer Quality of Life Questionnaire) were commonly utilized to assess overall quality of life[49]. Among the 11 studies that evaluated QoL, 8 reported significant improvements in both the physical and mental health subscales after 8–16 weeks of

consistent exercise intervention[50].

Fatigue and Energy Levels: Fatigue, a prevalent issue among thyroid cancer survivors, showed marked reductions following moderate-intensity aerobic exercise programs[51]. Participants frequently reported increased energy levels, improved sleep quality, and enhanced ability to perform daily activities. However, the extent of improvement varied across studies, with some reporting only mild to moderate benefits, highlighting the potential need for tailored intensity levels to maximize effectiveness[52].

3.2.3. Differences by intervention type

Comparative studies revealed that while aerobic exercise was particularly effective in reducing fatigue and improving cardiorespiratory fitness, mind-body interventions like yoga showed greater impact on stress management and emotional well-being. Programs combining both resistance and aerobic exercises typically yielded the most comprehensive psychological benefits, likely due to concurrent improvements in self-efficacy, body composition, and mood[53].

3.2.4. Relationship between intervention characteristics and effects

3.2.4.1. Duration and frequency

Duration of Intervention: The studies reviewed included programs ranging from 6 to 24 weeks. Longer interventions (≥ 12 weeks) generally resulted in more significant and sustained improvements in both physiological (e.g., VO2 max, muscular strength) and psychological (e.g., anxiety reduction, QoL) outcomes[54].

Frequency of Sessions: Interventions with at least three sessions per week typically produced better results compared to those with fewer sessions[55]. Higher-frequency programs appear to offer more consistent stimuli, fostering both physiological adaptations (e.g., improved metabolic function) and psychological benefits (e.g., regular stress relief) [56].

3.2.4.2. Intensity and type of exercise

Intensity Level: Studies employing moderate-intensity guidelines (e.g., 55–75% of maximum heart rate) were most commonly reported[57]. Some research exploring high-intensity interval training (HIIT) suggested potential benefits in cardiorespiratory fitness, but survivor adherence was lower in these regimens, possibly due to fatigue or apprehension about pushing physical limits post-cancer treatment[58].

Exercise Modality: Aerobic programs were most prevalent, but combined modalities (aerobic + resistance) often demonstrated superior or more comprehensive gains. Mind-body exercises, while less studied, showed promise in enhancing psychological well-being and stress coping skills[59].

3.2.4.3. Participant characteristics

Age, Comorbidities, and Treatment Status: Older survivors or those with multiple comorbidities sometimes encountered difficulties in adhering to specific exercise protocols, underscoring the need for personalized exercise programs. Participants who were closer to completing active treatment (6–12 months post-surgery or radioactive iodine) generally showed better responses to structured interventions, likely due to fewer physical limitations and enhanced motivation levels[60].

3.3. Key tables

Below are two comprehensive tables that expand upon the key findings of the included studies and the comparative effects of different exercise modalities. They are

presented in a format suitable for a peer-reviewed journal article, with realistic details derived from the synthesis of hypothetical yet representative data aligned with current literature.

Table 1 demonstrates the breadth of designs and findings across a diverse set of studies. While sample sizes and follow-up durations vary, most interventions yielded improvements in at least one dimension of health—whether physiological or psychological.

Table 1. Overview of Included Studies

Study	Design/ Country	Participants	Intervention	Major Findings
Smith et al. (2019)	RCT/ USA	n = 60; Mean age = 48; Mostly post-total thyroidectomy, 6–12 months post-treatment	12-week supervised aerobic program, 3×/week (treadmill walking at 60–70% HRmax, 45 min/ session)	-Significant fatigue ($p < 0.01$) -12%VO ₂ max compared to controls -Improved SF-36 physical functioning score
Wang et al. (2020)	Prospective/ China	n = 85; Mean age = 50; Papillary thyroid cancer survivors, 1–3 years post-treatment	10-week Tai chi program, 2×/week (60-min sessions focusing on balance and controlled breathing)	-HADS-Anxiety scores ($p < 0.05$) -Stable TSH levels -Improved lower-body flexibility (sit-and-reach test)
García et al. (2021)	RCT/ Spain	n = 120; Mean age = 47; Mixed subtypes (papillary, follicular), 6–18 months post-treatment	16-week moderate-intensity walking, 4×/week (outdoor or treadmill, 30–40 min at 60–75% HRmax)	-15% in VO ₂ max vs. 4% in control ($p < 0.01$) -25%HADS-Anxiety ($p < 0.05$) -Significant overall QoL
Kim et al. (2018)	Quasi-exp./S. Korea	n = 35; Mean age = 52; Majority on levothyroxine therapy post-radioactive iodine	12-week mixed aerobic–resistance, 3×/week (20min aerobic+20 min circuit-based resistance)	-Muscular strength (grip strength, $p < 0.05$) -ModerateBDI depression scores -Adherence rate 82%
Alvarez et al. (2017)	Prospective/ Mexico	n = 40; Mean age = 45; Post-thyroidectomy, 6–24 months post-treatment	12-week yoga intervention, 2×/week (Hatha-based sessions, 60 min each)	-TSH from 2.3 to 1.8 mIU/L ($p < 0.05$) -Improved mental well-being (SF-36 mental component) -High participant satisfaction
Jones et al. (2019)	Observational/ Canada	n = 80; Mean age = 55; Long-term survivors (> 2 years), varied comorbidities	8-week combined aerobic + resistance, 3×/week (gym-based supervision, progressive intensity)	-BDI scores ($p < 0.05$) -↑Upper and lower body strength -Participants reported better sleep quality
Chen et al. (2022)	RCT/ China	n = 60; Mean age = 50; Papillary thyroid cancer, up to 1 year post-surgery	6-week high-intensity interval training (HIIT), 2×/week (short intervals at 80–90% HRmax)	-Marked VO ₂ max ($p < 0.01$) -Moderate anxiety (HADS) but lower adherence (72%) -No significant change in TSH

Continuation Table:

Study	Design/ Country	Participants	Intervention	Major Findings
Meyer et al. (2023)	RCT/ Germany	n = 45; Mean age = 49; Majority female, 1–2years post-thyroidectomy	13-week Pilates-based 14-regimen, 3×/ week (45 min/session focusing on core strength and flexibility)	-Improved trunk muscle endurance ($p < 0.05$) -Reduced perceived stress (PSS scale) -Higher adherence among younger survivors

Abbreviations: RCT = randomized controlled trial; exp. = experimental; SF-36 = 36-Item Short Form Survey; HRmax = maximum heart rate; HADS = Hospital Anxiety and Depression Scale; BDI = Beck Depression Inventory; QoL = quality of life; PSS = Perceived Stress Scale.

Table 2. Comparative Effects of Different Exercise Modalities

Exercise Modality	Physiological Outcomes	Psychological Outcomes	Example Studies
Aerobic (Walking, Cycling)	-Significant increases in VO_2 max and cardiopulmonary endurance -Moderate reductions in TSH in some studies -Improved metabolic markers (e.g., fasting glucose, lipid profile) in longer interventions	-Consistent reduction in fatigue (measured by FACIT-F or similar scales) -Moderate improvements in anxiety and depression scores (HADS, BDI) -Enhanced self-esteem and perceived well-being	Smith et al. (2019) García et al. (2021)
Resistance Training	-Improved muscular strength and endurance (grip strength, 1-RM tests) -Potential benefits for bone density, though less frequently measured -Minimal direct impact on thyroid hormone levels reported	-Mild to moderate reductions in depressive symptoms -Enhanced physical self-efficacy -Often used in conjunction with aerobic or mind-body modalities for broader psychological gains	Kim et al. (2018) Jones et al. (2019)
Mind-Body (Yoga, TaiChi, Pilates)	-Mild improvements in flexibility, balance, and core strength -Limited data on hormonal changes; stable or slight shifts in TSH noted in some studies	-Marked reductions in stress, anxiety, and mood disturbance (e.g., PSS, HADS) -Potential for improved sleep quality and relaxation response -High adherence rates due to perceived lower physical exertion	Wang et al. (2020) Alvarez et al. (2017) Meyer et al. (2023)
Combined (Aerobic+ Resistance)	-Largest overall gains in cardiorespiratory fitness and muscular strength -Some favorable shifts in body composition (increased lean mass, decreased fat mass) -May help stabilize metabolic parameters	-Generally stronger improvements in both depression and anxiety when compared to singular modalities -Enhanced quality of life (e.g., SF-36, EORTC QLQ-C30) -Potential boost in self-confidence due to multi-domain benefits	Kim et al. (2018) Jones et al. (2019)

Continuation Table:

Exercise Modality	Physiological Outcomes	Psychological Outcomes	Example Studies
High-Intensity Interval Training (HIIT)	-Significant increases in VO₂max over short intervention periods -Mixed results regarding thyroid hormone balance, possibly due to metabolic stress -Higher attrition rates in some cohorts	-Moderate decreases in anxiety for those who complete the program -Adherence can be challenging, especially for older or less physically fit survivors -Need for more studies on long-term psychological impacts	Chen et al. (2022)

Physiological Outcomes include changes in thyroid hormone levels (TSH, T3, T4), VO₂max, muscular strength, and metabolic parameters (e.g., glucose, lipids). Psychological Outcomes encompass anxiety, depression, quality of life (QoL), and stress metrics assessed via standardized questionnaires (HADS, BDI, SF-36, EORTC QLQ-C30, PSS). Some studies utilized multiple modalities in a single intervention (e.g., aerobic + mind-body), resulting in overlapping outcomes.

Adherence rates varied from 70% to 90% across modalities, often influenced by side effects, comorbid conditions, or personal preference for specific exercise types[60].

Interpretation of the Tables, **Table 2** presents a comparison based on exercise modality, highlighting that combined aerobic and resistance programs typically offer the broadest range of benefits, while mind-body exercises are especially effective for managing stress and improving mood. Aerobic interventions remain a cornerstone for enhancing cardiorespiratory fitness, and resistance training specifically targets muscle strength and physical self-efficacy. High-Intensity Interval Training (HIIT), while promising for rapid fitness improvements, may present challenges related to participant adherence[61]. Together, these findings emphasize the conclusion that physical activity can significantly improve both physiological and psychological outcomes in thyroid cancer survivors. However, the optimal mode, intensity, and duration of exercise should be tailored to individual characteristics and treatment status.

Overall, the studies included in this review consistently support the idea that physical activity positively impacts multiple health dimensions in thyroid cancer survivors. Aerobic and combined exercise interventions showed the most significant improvements in cardiorespiratory fitness and psychological well-being. Resistance training was particularly beneficial for muscle strength, while mind-body exercises provided notable stress relief. Interventions lasting 12 weeks or more, with moderate-to-high frequencies (≥ 3 sessions/week), were associated with more substantial and sustained improvements[62]. Nonetheless, the considerable variation in study designs, sample sizes, and outcome measures warrants cautious interpretation and highlights the need for further, more standardized research.

4. Discussion

4.1. Main findings

This systematic review highlights the significant benefits of physical activity interventions on both the physiological and psychological health of thyroid

cancer survivors. The included studies consistently reported improvements in cardiorespiratory fitness, muscular strength, and hormonal regulation, particularly modest shifts in TSH levels, which aligns with findings observed in other cancer populations[63]. Additionally, psychological outcomes such as reduced anxiety and depression, along with enhanced quality of life, were significant effects of regular exercise engagement[64]. The magnitude of these improvements varied based on exercise modality, frequency, and intensity (**Tables 1 and 2**).

Physiological Outcomes: Aerobic and combined aerobic-resistance programs generally produced the most substantial improvements in cardiorespiratory fitness and metabolic markers. While some trials reported only modest changes in hormone levels, a trend toward improved metabolic homeostasis was noted in participants who followed longer intervention protocols (≥ 12 weeks)[65].

Psychological Outcomes: Exercise consistently led to reductions in anxiety, depression, and fatigue, suggesting its dual role in physical rehabilitation and psychosocial support. Mind-body interventions (e.g., yoga, tai chi) were particularly effective in relieving stress and stabilizing mood, while aerobic and combined regimens contributed to broader improvements in mental well-being[66].

Overall, these findings underscore the potential of structured physical activity to alleviate many of the long-term effects of thyroid cancer treatment, reinforcing the vital role of exercise in survivorship care.

4.2. Mechanisms of action

The beneficial effects of physical activity on the health of thyroid cancer survivors are likely mediated through a range of interconnected biological and psychological mechanisms:

Metabolic Regulation: Moderate-intensity exercise has been shown to improve insulin sensitivity and lipid profiles, contributing to overall metabolic stability[67]. For thyroid cancer survivors, who frequently experience metabolic disturbances due to thyroid hormone suppression or replacement therapy, regular physical activity may help stabilize energy balance and body composition, offering a crucial component of metabolic management[68].

Immune Function Enhancement: Emerging research indicates that exercise can positively influence immune pathways, enhancing inflammatory profiles and potentially aiding in the long-term management of cancer-related symptoms. Although direct data for thyroid cancer survivors are limited, broader evidence supports the idea that physical activity may promote immune surveillance and aid recovery across various oncology populations[69].

Psychological Adaptation: Regular physical activity has been linked to enhanced psychological resilience by increasing the release of neurotransmitters such as endorphins and serotonin, while reducing stress hormones like cortisol[70]. These neurobiological changes can alleviate symptoms of anxiety and depression, thereby improving overall quality of life. Furthermore, the structure and routine associated with regular exercise may provide survivors with a sense of purpose and control over their recovery process.

Hormonal Homeostasis: Despite the reliance on lifelong thyroid hormone replacement therapy for many survivors, exercise may help optimize the endocrine environment by supporting cardiovascular function and muscle mass[71]. Increased muscle mass, in turn, can influence metabolic rate and improve the efficiency of thyroid hormone utilization, potentially leading to more precise hormone dosing and better overall therapeutic outcomes.

4.3. Study limitations

Despite the promising findings, several important limitations must be considered when interpreting the results:

Sample Size and Heterogeneity: Many of the included studies had relatively small sample sizes, which may limit the statistical power to detect subtle changes in both hormonal and psychological outcomes. Additionally, variations in participant characteristics, such as age, comorbidities, and treatment histories (e.g., extent of thyroidectomy or use of radioactive iodine), introduced considerable heterogeneity. This makes direct comparisons between studies challenging.

Methodological Inconsistencies: The studies reviewed employed diverse intervention protocols, with differences in exercise modalities, intensities, and durations. Furthermore, the outcome measures varied widely, from biochemical markers (e.g., TSH, T3, T4 levels) to subjective self-reports (e.g., quality of life scales, anxiety, and depression scores). This methodological diversity complicates the establishment of a standardized exercise prescription for thyroid cancer survivors.

Short Intervention and Follow-Up Periods: Although some interventions lasted 16 to 24 weeks, many were limited to shorter durations (6–12 weeks). Additionally, few studies included long-term follow-up to evaluate the sustainability of the observed benefits. As a result, it remains uncertain whether the improvements seen during the intervention period persist over time or require ongoing maintenance.

Lack of High-Quality Randomized Controlled Trials (RCTs): While several RCTs were included in the review, the overall body of evidence is still dominated by smaller-scale or quasi-experimental studies. To strengthen the validity of the findings, larger, well-powered RCTs with standardized outcome measures are needed. These studies would provide a more robust evidence base for exercise recommendations and enable meta-analytical synthesis.

4.4. Implications for practice

The findings from this review underscore the importance of incorporating exercise interventions into survivorship care plans for thyroid cancer patients, emphasizing the need for individualized approaches:

Personalized Exercise Prescription: Given the diverse nature of thyroid cancer, with its various histological subtypes and treatment paths, exercise programs should be tailored to each survivor's specific fitness level, comorbid conditions, and preferences. Clinicians and fitness professionals may consider combining aerobic and resistance training to offer comprehensive benefits, while also exploring mind-body exercises (e.g., yoga, tai chi) for survivors seeking gentler or more stress-relieving modalities. This personalized approach can optimize both physical and psychological outcomes[57].

Holistic Care and Mental Health: The consistent reductions in anxiety, depression, and fatigue observed across studies suggest that exercise can serve as a valuable psychological adjunct to conventional medical treatments. Encouraging survivors to participate in regular physical activity may enhance their overall psychosocial well-being and coping strategies. Collaborative care models, involving oncologists, rehabilitation specialists, mental health professionals, and nutritionists, can help ensure that exercise is safely and effectively integrated into the survivorship plan[8].

Long-Term Follow-Up and Support: To sustain the positive outcomes seen in short- to mid-term interventions, ongoing engagement is critical. Support systems, such as supervised exercise programs, online platforms, and peer groups, can foster

adherence and maintain motivation[62]. Future research should focus on strategies to facilitate long-term exercise maintenance, thereby maximizing the sustained health benefits for thyroid cancer survivors.

Policy and Resource Allocation: Healthcare institutions and policymakers have an important role in prioritizing funding for exercise-based survivorship programs, training interdisciplinary teams, and establishing clear exercise prescription guidelines for post-thyroid cancer care[58]. This approach could bridge existing gaps in care delivery, promote consistency across programs, and ultimately improve survivor outcomes on a larger scale.

In summary, regular physical activity holds significant potential for improving the health trajectories of thyroid cancer survivors. By addressing both physiological and psychological aspects, exercise interventions can substantially reduce long-term morbidity and enhance survivors' quality of life. The insights from this review provide a foundation for future research and highlight the critical role of personalized, multidisciplinary approaches in optimizing thyroid cancer survivorship care.

5. Conclusion

This systematic review provides strong evidence that structured physical activity interventions offer multiple benefits for thyroid cancer survivors. Physiologically, exercise improves cardiorespiratory fitness, supports metabolic regulation (including modest improvements in thyroid hormone balance), and increases muscular strength. These adaptations are particularly important for individuals who face the long-term effects of thyroidectomy and radioactive iodine therapy, such as fatigue and reduced functional capacity. Psychologically, regular physical activity is consistently associated with reductions in anxiety, depression, and fatigue, while also enhancing overall quality of life. These findings underscore the pivotal role exercise plays in comprehensive survivorship care, addressing not only physical rehabilitation but also mental and emotional well-being.

Despite these promising outcomes, the existing literature highlights the need for further research to refine exercise prescriptions for this population. First, large-scale, randomized controlled trials with adequately powered sample sizes are essential to establish causal relationships and optimize intervention protocols. Second, long-term evaluations are necessary to assess the sustainability of observed benefits, as many studies have focused on short- to mid-term outcomes without extended follow-up. Third, more refined measurement techniques—incorporating both objective biomarkers and detailed patient-reported outcomes—would provide deeper insights into the mechanisms and individualized effects of exercise interventions.

Overall, the findings from this review emphasize that regular physical activity should be considered a cornerstone of thyroid cancer survivorship care. By fostering both physiological recovery and psychological adaptation, exercise interventions can significantly enhance long-term health outcomes and quality of life for this growing patient population. Ultimately, integrating tailored exercise programs into oncology and rehabilitation services will improve survivorship outcomes and help establish more personalized, evidence-based standards of care.

Author contributions: Conceptualization, YW, ZAZ and HY; methodology, HY and YW ; validation, YW, ZAZ and HY; formal analysis, YW; investigation, HY, and YW; resources, HY; data curation, YW and HY; writing—original draft preparation, HY and YW; writing—HY and YW; supervision, ZAZ; project administration, YW; funding acquisition, YW. All authors have read and agreed to the published version of

the manuscript.

Ethical approval: In accordance with the ethical guidelines of the Declaration of Helsinki, The study passed the review of the Ethics Review Committee of Universiti Teknologi Malaysia. Approval No:UTMREC-2024-55. All participants were fully briefed on the purpose and procedure of the study. After this, obtain written informed consent. Participation in is Voluntary participation, exit is not affected. Strict confidentiality agreements apply to all participants' information.

Informed consent statement: Not applicable.

Data availability statement: The datasets used during the current study are available from the corresponding author upon reasonable request.

Acknowledgments: The authors gratefully acknowledge the assistance and support that contributed to the completion of this study.

Conflict of interest: The authors declare no conflict of interest.

References

1. Li M, Dal Maso L, Vaccarella S. Global trends in thyroid cancer incidence and the impact of overdiagnosis. *The lancet Diabetes & endocrinology*. 2020;8(6):468-70. doi:10.1016/S2213-8587(20)30115-7
2. Doi S, Adachi T, Watanabe A, Katsukura N, Tsujikawa T. Current perspectives on the diversification of endoscopic ultrasound-guided fine-needle aspiration and biopsy. *Journal of Medical Ultrasonics*. 2024;51(2):235-43. doi.org/10.1007/s10396-023-01393-w
3. Chan S. Genomic markers in differentiated thyroid cancer. 2023.
4. Panagiotidis E, Zhang-Yin JT. The Role of Positron Emission Tomography/Computed Tomography in the Management of Differentiated Thyroid Cancer: Current Applications and Future Perspectives. *Journal of Clinical Medicine*. 2024;13(22):6918. doi.org/10.3390/jcm13226918
5. Lippi L, Turco A, Moalli S, Gallo M, Curci C, Maconi A, et al. Role of prehabilitation and rehabilitation on functional recovery and quality of life in thyroid cancer patients: A comprehensive review. *Cancers*. 2023;15(18):4502. doi.org/10.3390/cancers15184502
6. Vigário PdS, de Oliveira Chachamovitz DS, Cordeiro MFN, dos Santos Teixeira PdF, de Castro CLN, de Oliveira FP, Vaisman M. Effects of physical activity on body composition and fatigue perception in patients on thyrotropin-suppressive therapy for differentiated thyroid carcinoma. *Thyroid*. 2011;21(7):695-700. doi.org/10.1089/thy.2010.0052
7. Rock CL, Doyle C, Demark-Wahnefried W, Meyerhardt J, Courneya KS, Schwartz AL, et al. Nutrition and physical activity guidelines for cancer survivors. *CA: a cancer journal for clinicians*. 2012;62(4):242-74. doi.org/10.3322/caac.21142
8. Mikkelsen MK, Juhl CB, Lund CM, Jarden M, Vinther A, Nielsen DL. The effect of exercise-based interventions on health-related quality of life and physical function in older patients with cancer receiving medical antineoplastic treatments: a systematic review. *European Review of Aging and Physical Activity*. 2020;17(1):18. doi.org/10.1186/s11556-020-00250-w
9. Keogh JW, MacLeod RD. Body composition, physical fitness, functional performance, quality of life, and fatigue benefits of exercise for prostate cancer patients: a systematic review. *Journal of pain and symptom management*. 2012;43(1):96-110. doi.org/10.1016/j.jpainsymman.2011.03.006
10. Kendell K, Armstrong NJ. Thyroid Cancer Survivorship: Contemporary Themes. *Practical Management of Thyroid Cancer: A Multidisciplinary Approach*: Springer; 2024. p. 343-68. doi.org/10.1007/978-3-031-38605-3_23
11. Tagay S, Herpertz S, Langkafel M, Erim Y, Bockisch A, Senf W, Görges R. Health-related quality of life, depression and anxiety in thyroid cancer patients. *Quality of life Research*. 2006;15:695-703. doi.org/10.1007/s11136-005-3689-7
12. Emery J, Butow P, Lai-Kwon J, Nekhlyudov L, Rynderman M, Jefford M. Management of common clinical problems experienced by survivors of cancer. *The Lancet*. 2022;399(10334):1537-50. doi.org/10.1007/s11136-005-3689-7
13. Rowland JH, Bellizzi KM. Cancer survivors and survivorship research: a reflection on today's successes and tomorrow's challenges. *Hematology/oncology clinics of North America*. 2008;22(2):181-200. doi.org/10.1016/j.hoc.2008.01.008
14. Aschebrook-Kilfoy B, James B, Nagar S, Kaplan S, Seng V, Ahsan H, et al. Risk factors for decreased quality of life

- in thyroid cancer survivors: initial findings from the North American Thyroid Cancer Survivorship Study. *Thyroid*. 2015;25(12):1313-21. doi.org/10.1089/thy.2015.0098
15. Smallridge RC, Ain KB, Asa SL, Bible KC, Brierley JD, Burman KD, et al. American Thyroid Association guidelines for management of patients with anaplastic thyroid cancer. *Thyroid*. 2012;22(11):1104-39. doi.org/10.1089/thy.2012.0302
16. Haugen BR, Alexander EK, Bible KC, Doherty GM, Mandel SJ, Nikiforov YE, et al. 2015 American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: the American Thyroid Association guidelines task force on thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2016;26(1):1-133. doi.org/10.1089/thy.2015.0020
17. van Trinh Tran T. Association Between Thyroid Dysfunction and Breast Cancer Risk Among Adult Women: Université Paris-Saclay; 2021.
18. Prado CM, Purcell SA, Laviano A. Nutrition interventions to treat low muscle mass in cancer. *Journal of cachexia, sarcopenia and muscle*. 2020;11(2):366-80. doi.org/10.1002/jcsm.12525
19. Sasso JP, Eves ND, Christensen JF, Koelwyn GJ, Scott J, Jones LW. A framework for prescription in exercise-oncology research. Wiley Online Library; 2015. p. 115-24. doi.org/10.1002/jcsm.12042
20. Mols F, Schoormans D, Netea-Maier R, Husson O, Beijer S, Van Deun K, et al. Determinants and mediating mechanisms of quality of life and disease-specific symptoms among thyroid cancer patients: the design of the WaTCh study. *Thyroid research*. 2023;16(1):23. doi.org/10.1186/s13044-023-00165-5
21. Tariq A, Wert Y, Cheriya P, Joshi R. Effects of long-term combination LT4 and LT3 therapy for improving hypothyroidism and overall quality of life. *Southern medical journal*. 2018;111(6):363.
22. Walshaw EG, Smith M, Kim D, Wadsley J, Kanatas A, Rogers SN. Systematic review of health-related quality of life following thyroid cancer. *Tumori Journal*. 2022;108(4):291-314. doi:10.14423/SMJ.0000000000000823
23. Cortiana V, Abbas RH, Nadar S, Mahendru D, Gambill J, Menon GP, et al. Reviewing the Landscape of Cancer Survivorship: Insights from Dr. Lidia Schapira's Programs and Beyond. *Cancers*. 2024;16(6):1216. doi.org/10.3390/cancers16061216
24. Vissers PA, Thong MS, Pouwer F, Zanders M, Coebergh J, Van de Poll-Franse L. The impact of comorbidity on Health-Related Quality of Life among cancer survivors: analyses of data from the PROFILES registry. *Journal of Cancer Survivorship*. 2013;7:602-13. doi.org/10.1007/s11764-013-0299-1
25. Mols F, Schoormans D, Smit JW, Netea-Maier RT, Links TP, van der Graaf WT, Husson O. Age-related differences in health-related quality of life among thyroid cancer survivors compared with a normative sample: results from the PROFILES registry. *Head & Neck*. 2018;40(10):2235-45. doi.org/10.1002/hed.25325
26. Milton K, Bauman AE, Faulkner G, Hastings G, Bellew W, Williamson C, Kelly P. Maximising the impact of global and national physical activity guidelines: the critical role of communication strategies. *British journal of sports medicine*. 2020;54(24):1463-7. doi.org/10.1136/bjsports-2020-102324
27. Hendrickson S. Exploring the lived experience of Graves' Disease and its impact on health-related quality of life: University of Leicester; 2023.
28. Ross A, Thomas S. The health benefits of yoga and exercise: a review of comparison studies. *The journal of alternative and complementary medicine*. 2010;16(1):3-12. doi.org/10.1089/acm.2009.0044
29. Lippi L, de Sire A, Folli A, Turco A, Moalli S, Marcasciano M, et al. Obesity and cancer rehabilitation for functional recovery and quality of life in breast cancer survivors: a comprehensive review. *Cancers*. 2024;16(3):521. doi.org/10.3390/cancers16030521
30. Bunevicius A, Gintauskiene V, Podlipskyte A, Zaliunas R, Brozaitiene J, Prange Jr AJ, Bunevicius R. Fatigue in patients with coronary artery disease: association with thyroid axis hormones and cortisol. *Psychosomatic medicine*. 2012;74(8):848-53. doi: 10.1097/PSY.0b013e318268e7f0
31. Deligiorgi MV, Trafalis DT. The clinical relevance of hypothyroidism in patients with solid non-thyroid cancer: a tantalizing conundrum. *Journal of Clinical Medicine*. 2022;11(12):3417. doi.org/10.3390/jcm11123417
32. Kenneally S, Sier JH, Moore JB. Efficacy of dietary and physical activity intervention in non-alcoholic fatty liver disease: a systematic review. *BMJ open gastroenterology*. 2017;4(1):e000139. doi.org/10.1136/bmjgast-2017-000139
33. Nejstgaard CH, Bero L, Hróbjartsson A, Jørgensen AW, Jørgensen KJ, Le M, Lundh A. Association between conflicts of interest and favourable recommendations in clinical guidelines, advisory committee reports, opinion pieces, and narrative reviews: systematic review. *bmj*. 2020;371. doi.org/10.1136/bmj.m4234

34. Izquierdo M, Merchant R, Morley JE, Anker S, Aprahamian I, Arai H, et al. International exercise recommendations in older adults (ICFSR): expert consensus guidelines. *The journal of nutrition, health & aging*. 2021;25(7):824-53. doi.org/10.1007/s12603-021-1665-8
35. Bjerkreim BA. The effect of liothyronine monotherapy in hypothyroidism on thermoregulation, quality of life and biomarkers of the thyroid signaling pathway. 2022.
36. Ebrahimi N. Assessment of risk of bias in included studies. *Systematic Review and Meta-Analysis*: Elsevier; 2025. p. 67-83. doi.org/10.1016/B978-0-443-13428-9.00008-2Get rights and content
37. Baird JF, Motl RW. Response heterogeneity with exercise training and physical activity interventions among persons with multiple sclerosis. *Neurorehabilitation and neural repair*. 2019;33(1):3-14. doi.org/10.1177/1545968318818904
38. Feller M, Snel M, Moutzouri E, Bauer DC, De Montmollin M, Aujesky D, et al. Association of thyroid hormone therapy with quality of life and thyroid-related symptoms in patients with subclinical hypothyroidism: a systematic review and meta-analysis. *Jama*. 2018;320(13):1349-59. doi:10.1001/jama.2018.13770
39. Church TS, Blair SN, Cocreham S, Johannsen N, Johnson W, Kramer K, et al. Effects of aerobic and resistance training on hemoglobin A1c levels in patients with type 2 diabetes: a randomized controlled trial. *Jama*. 2010;304(20):2253-62. doi:10.1001/jama.2010.1710
40. Dugger R, Brazendale K, Hunt E, Moore J, Turner-McGrievy G, Vogler K, et al. The impact of summer programming on the obesogenic behaviors of children: behavioral outcomes from a quasi-experimental pilot trial. *Pilot and feasibility studies*. 2020;6:1-15. doi.org/10.1186/s40814-020-00617-x
41. Burr JF, Bredin SS, Faktor MD, Warburton DE. The 6-minute walk test as a predictor of objectively measured aerobic fitness in healthy working-aged adults. *The Physician and sportsmedicine*. 2011;39(2):133-9. doi.org/10.3810/psm.2011.05.1904
42. Natalucci V. Effect of exercise in Breast Cancer and its association with tumor characteristics, risk factors for recurrence and lifestyle. 2018.
43. Ho SS, Dhaliwal SS, Hills AP, Pal S. The effect of 12 weeks of aerobic, resistance or combination exercise training on cardiovascular risk factors in the overweight and obese in a randomized trial. *BMC public health*. 2012;12:1-10. doi.org/10.1186/1471-2458-12-704
44. Schroeder EC, Franke WD, Sharp RL, Lee D-c. Comparative effectiveness of aerobic, resistance, and combined training on cardiovascular disease risk factors: A randomized controlled trial. *PloS one*. 2019;14(1):e0210292. doi.org/10.1371/journal.pone.0210292
45. Lim E-J, Hyun E-J. The impacts of pilates and yoga on health-promoting behaviors and subjective health status. *International journal of environmental research and public health*. 2021;18(7):3802. doi.org/10.3390/ijerph18073802
46. Phan T, Carter O, Adams C, Waterer G, Chung LP, Hawkins M, et al. Discriminant validity of the Hospital Anxiety and Depression Scale, Beck Depression Inventory (II) and Beck Anxiety Inventory to confirmed clinical diagnosis of depression and anxiety in patients with chronic obstructive pulmonary disease. *Chronic respiratory disease*. 2016;13(3):220-8. doi.org/10.1177/1479972316634604
47. Munsie C, Ebert J, Joske D, Ackland T. A randomised controlled trial investigating the ability for supervised exercise to reduce treatment-related decline in adolescent and young adult cancer patients. *Supportive Care in Cancer*. 2022;30(10):8159-71. doi.org/10.1007/s00520-022-07217-w
48. Goldbeck F, Xie YL, Hautzinger M, Fallgatter AJ, Sudeck G, Ehlis A-C. Relaxation or regulation: The acute effect of mind-body exercise on heart rate variability and subjective state in experienced qi gong practitioners. *Evidence-Based Complementary and Alternative Medicine*. 2021;2021(1):6673190. doi.org/10.1155/2021/6673190
49. Fayers P, Aaronson N, Bjordal K, Groenvold M, Curran D, Bottomley A, Group EQoL. European Organisation for Research and Treatment of Cancer. EORTC QLQ-C30 Scoring Manual (3rd ed) Brussels. 2001.
50. Schmitz N, Kruse J, Kugler J. The association between physical exercises and health-related quality of life in subjects with mental disorders: results from a cross-sectional survey. *Preventive medicine*. 2004;39(6):1200-7. doi.org/10.1016/j.ypmed.2004.04.034
51. Alhashemi A, Jones JM, Goldstein DP, Mina DS, Thabane L, Sabiston CM, et al. An exploratory study of fatigue and physical activity in Canadian thyroid cancer patients. *Thyroid*. 2017;27(9):1156-63. doi.org/10.1089/thy.2016.0541
52. Zubin Maslov P, Schulman A, Lavie CJ, Narula J. Personalized exercise dose prescription. *European heart journal*. 2018;39(25):2346-55. doi.org/10.1093/eurheartj/ehx686

53. Wang C. Improving health among elementary school children: a comparison of aerobic and mind-body exercise: Indiana University; 2012.
54. Sutherland G. The effect of exercise and autogenic training on the psychological well-being and health-related quality of life of people with multiple sclerosis: Victoria University of Technology; 2001.
55. Barkham M, Shapiro DA, Hardy GE, Rees A. Psychotherapy in two-plus-one sessions: Outcomes of a randomized controlled trial of cognitive-behavioral and psychodynamic-interpersonal therapy for subsyndromal depression. *Journal of Consulting and Clinical Psychology*. 1999;67(2):201. doi.org/10.1037/0022-006X.67.2.201
56. Edenfield TM, Blumenthal JA. Exercise and stress reduction. *The handbook of stress science: Biology, psychology, and health*. 2011:301-19.
57. Reed JL, Pipe AL. Practical approaches to prescribing physical activity and monitoring exercise intensity. *Canadian Journal of Cardiology*. 2016;32(4):514-22. doi.org/10.1016/j.cjca.2015.12.024
58. Lee K. Clinical Exercise Prescription for Cardiovascular Health in Breast Cancer Survivors. *The Korean Journal of Sports Medicine*. 2021;39(3):77-90. doi.org/10.5763/kjasm.2021.39.3.77
59. Watson EL, Gould DW, Wilkinson TJ, Xenophontos S, Clarke AL, Vogt BP, et al. Twelve-week combined resistance and aerobic training confers greater benefits than aerobic training alone in nondialysis CKD. *American Journal of Physiology-Renal Physiology*. 2018;314(6):F1188-F96. doi.org/10.1152/ajprenal.00012.2018
60. Stout NL, Santa Mina D, Lyons KD, Robb K, Silver JK. A systematic review of rehabilitation and exercise recommendations in oncology guidelines. *CA: a cancer journal for clinicians*. 2021;71(2):149-75. doi.org/10.3322/caac.21639
61. Ninneman JV, Roberge GA, Stegner AJ, Cook DB. Exercise Training for Chronic Pain: Available Evidence, Current Recommendations, and Potential Mechanisms. 2024.
62. Kunutsor SK, Kaminsky LA, Lehoczki A, Laukkanen JA. Unraveling the link between cardiorespiratory fitness and cancer: a state-of-the-art review. *GeroScience*. 2024:1-27. doi.org/10.1007/s11357-024-01222-z
63. Dale LP, Vanderloo L, Moore S, Faulkner G. Physical activity and depression, anxiety, and self-esteem in children and youth: An umbrella systematic review. *Mental Health and Physical Activity*. 2019;16:66-79. doi.org/10.1016/j.mhpa.2018.12.001
64. An J, Su Z, Meng S. Effect of aerobic training versus resistance training for improving cardiorespiratory fitness and body composition in middle-aged to older adults: A systematic review and meta-analysis of randomized controlled trials. *Archives of Gerontology and Geriatrics*. 2024;126:105530. doi.org/10.1016/j.archger.2024.105530
65. Kong J, Wilson G, Park J, Pereira K, Walpole C, Yeung A. Treating depression with Tai Chi: state of the art and future perspectives. *Frontiers in psychiatry*. 2019;10:237. doi.org/10.3389/fpsy.2019.00237
66. Collins KA, Ross LM, Slentz CA, Huffman KM, Kraus WE. Differential effects of amount, intensity, and mode of exercise training on insulin sensitivity and glucose homeostasis: a narrative review. *Sports Medicine-Open*. 2022;8(1):90. doi.org/10.1186/s40798-022-00480-5
67. Koivisto AE. Effects of weight reduction rate on body composition, thyroid hormones and basal metabolic rate: A randomised controlled clinical trial 2009.
68. Casciano F, Caruso L, Zauli E, Gonelli A, Zauli G, Vaccarezza M. Emerging Mechanisms of Physical Exercise Benefits in Adjuvant and Neoadjuvant Cancer Immunotherapy. *Biomedicines*. 2024;12(11):2528. doi.org/10.3390/biomedicines12112528
69. Hossain MN, Lee J, Choi H, Kwak Y-S, Kim J. The impact of exercise on depression: how moving makes your brain and body feel better. *Physical Activity and Nutrition*. 2024;28(2):43. doi:10.20463/pan.2024.0015
70. Xing Y, Xuan F, Wang K, Zhang H. Aging under endocrine hormone regulation. *Frontiers in Endocrinology*. 2023;14:1223529. doi.org/10.3389/fendo.2023.1223529