

Review

Educational applications of aerobic exercise in psycho-oncology: Teaching mind–body resilience through movement

Jin Xu[#], Wenjing Guo[#], Jianmin Dai^{*}

Kyungnam University, 7 Kyungnamdaehak-ro, Masanhappo-gu, Changwon-si, Gyeongsangnam-do 51276, South Korea

*** Corresponding author:** Jianmin Dai, daijianmin950210@126.com

[#] These authors are co-first authors and contributed equally to this manuscript.

CITATION

Xu J, Guo W, Dai J. Educational applications of aerobic exercise in psycho-oncology: Teaching mind–body resilience through movement. *Psycho-Oncologie*. 2025; 19(4): 5564.
<https://doi.org/10.18282/po5564>

ARTICLE INFO

Received: 15 September 2025
Revised: 18 October 2025
Accepted: 12 November 2025
Available online: 12 December 2025

COPYRIGHT



Copyright © 2025 by author(s).
Psycho-Oncologie is published by PiscoMed Publishing Pte. Ltd. This work is licensed under the Creative Commons Attribution (CC BY) license.
<https://creativecommons.org/licenses/by/4.0/>

Abstract: Cancer represents not only a biomedical condition but also a profound psychological and existential experience that challenges patients' sense of agency and well-being. Within psycho-oncology, aerobic exercise has emerged as a powerful mind–body intervention that extends beyond physical rehabilitation to serve as an educational medium for cultivating resilience. This narrative review synthesizes theoretical, empirical, and pedagogical evidence to explore how structured aerobic activity can teach self-regulation, emotional awareness, and adaptive coping in cancer care. Drawing on biopsychosocial and neuropsychobiological models, the review demonstrates that rhythmic aerobic movement enhances mood, cognition, and physiological coherence while fostering reflective learning through embodied experience. Integrating these effects into psycho-oncological education transforms exercise into a cyclical process of physiological activation, emotional insight, reflection, and personal transformation. Implementing such movement-based pedagogy requires interdisciplinary collaboration, culturally sensitive adaptation, and reflective facilitation, yet it holds significant promise for both patients and healthcare professionals. Ultimately, teaching resilience through movement unites therapy and education, embodying the biopsychosocial principles of psycho-oncology and redefining recovery as a dynamic, experiential process through which individuals learn to move with awareness, strength, and purpose toward holistic well-being.

Keywords: aerobic exercise; psycho-oncology; resilience; mind–body education; cancer survivorship; embodied learning

1. Introduction

Cancer is both a significant biomedical challenge and a profound psychological and existential crisis for patients. This duality is evident in the myriad struggles faced by individuals undergoing cancer treatment, including physical discomfort, emotional distress, and a disruptive shift in one's sense of agency and purpose. Cancer patients often suffer from anxiety, fear of recurrence, social withdrawal, and a pervasive sense of helplessness stemming from their condition and treatment regimens [1,2]. The emergence of psycho-oncology as a discipline seeks to address these complexities by advocating for a holistic approach that integrates psychological and physical dimensions of care, emphasizing that recovery is fundamentally a re-integration of mind, body, and emotion [3,4].

Central to the psycho-oncological framework is the concept of resilience, defined as the capacity to adapt and recover from adversity. Resilience is not a static trait but a dynamic process that can be fostered through various means, including educational and experiential interventions [3,4]. Research indicates that resilience can be significantly enhanced through structured programs that incorporate both physical

activity and psychological support, effectively mitigating the emotional distress experienced by cancer patients [4,5]. Aerobic exercise has gained traction as a core component of this integrative approach, transitioning from a mere adjunct to a pivotal psycho-physiological intervention that positively influences mood, cognitive performance, and self-perception [6,7].

Engaging in aerobic activities such as walking, cycling, or dance promotes a host of neurobiological benefits, including improved stress response, enhanced cerebral blood flow, and optimized immune function, resulting in measurable emotional and cognitive enhancements [8,9]. The rhythmic nature of these exercises contributes to neurotrophic and endorphin release, fostering greater emotional regulation and cognitive clarity tools that are essential for counteracting the helplessness often associated with cancer experiences [8,9]. Consequently, aerobic exercise facilitates emotional balance and empowers patients with a renewed sense of self-efficacy, allowing them to navigate their treatment journey with a greater sense of control [2,9,10].

While the therapeutic benefits of exercise are increasingly recognized, its potential as an educational tool remains underexplored. Educational frameworks in health care emphasize the importance of integrative learning, where knowledge acquisition is deepened through embodied experiences [11]. By embedding movement and exercise into psycho-oncological education, educators can demonstrate the physiological connections between bodily regulation and emotional states, enabling both healthcare professionals and patients to learn self-management skills and resilience in practice [12,13]. Movement education, particularly in the context of psycho-oncology, transforms physical activity into a pedagogical act, where participants experience firsthand how physical regulation can enhance emotional resilience [1,11,13].

Furthermore, integrating aerobic exercise into educational frameworks can facilitate meaningful communication regarding emotional competence and self-management among medical students and healthcare providers. This underscores the necessity of adopting a “hidden curriculum” that cultivates emotional intelligence alongside medical training [14]. Such educational constructs support future practitioners in developing empathy and equip them with practical strategies to address the psychological dimensions of patient care, particularly in high-stress settings like oncology [14].

Despite the recognized benefits of movement-based interventions, this field remains underdeveloped in systematic research. Current psycho-oncological programs primarily focus on outcome measurements rather than on the educational methodologies that can cultivate resilience through movement [4]. There is a pressing need for studies that bridge neuroscience, psychology, and pedagogy to create coherent models for integrating movement as a core educational approach in psycho-oncology. This narrative review advocates for further exploration into how aerobic exercise can function as a curriculum of resilience, granting patients and healthcare providers the tools necessary for embodied learning and recovery [1,2,9].

In conclusion, fostering resilience through movement represents a transformative goal for psycho-oncology, reconceptualizing physical activity as a collaborative

learning journey. This approach emphasizes the interplay of bodily rhythm with emotional well-being, reinforcing the notion that the mind and body are inseparable in the healing process. Thus, the educational applications of aerobic exercise pave the way for an innovative pathway that integrates therapy, pedagogy, and personal growth in cancer care, advocating a holistic approach to health where learning and healing coexist [2,13,14].

2. Theoretical foundations

2.1. The biopsychosocial framework in psycho-oncology

Psycho-oncology has significantly evolved, shifting the understanding of cancer from a solely biological phenomenon to a multifaceted interaction of psychological, behavioral, and social dimensions. This evolution is underpinned by the biopsychosocial model introduced by Engel, which posits that health and disease result from a dynamic interplay between biological, psychological, and social factors rather than merely from cellular pathological processes [15]. This paradigm emphasizes the necessity of addressing emotional distress, cognitive appraisals, and social contexts, all of which critically shape a patient's disease experience, adherence to treatment, and overall quality of life [16].

Within this framework, resilience is conceptualized as a dynamic process rather than a fixed trait, defining the individual's ability to adaptively respond to adversity [17]. The biopsychosocial model recognizes that resilience encompasses an interplay of biological (e.g., immune function), psychological (e.g., emotional regulation), and social (e.g., support networks) factors and highlights how these subsystems affect one another [18]. For instance, physiological stability enhances emotional regulation, while robust social connections foster meaning-making and hope. Conversely, chronic stress or social isolation can lead to dysregulation of neuroendocrine and immune systems, heightening vulnerability to both psychological and physical deterioration [19].

In educational contexts, this model serves both scientific and pedagogical functions, teaching healthcare learners to approach resilience as a systemic competence. It empowers educators to illustrate how techniques like cognitive reframing, emotional expression, and physical self-regulation function as interconnected pathways to adaptation, further reinforcing the need for integrating aerobic exercise within psycho-oncological training as a tangible representation of the biopsychosocial interplay [20]. By actively engaging in aerobic activity, individuals experience firsthand the dynamics of their biopsychosocial systems during movement, thereby embodying the lessons of resilience [18].

2.2. Exercise and neuropsychobiological resilience

Aerobic exercise has been shown to exert regulatory effects aligning with resilience mechanisms at the neuropsychobiological level. Regular physical activity activates the neuroendocrine and immune systems that are pivotal for stress management and adaptive responses [21]. Key neurobiological substrates, such as brain-derived neurotrophic factor (BDNF) and endorphins, are released during

exercise, promoting neuroplasticity and mood regulation, while simultaneous reductions in cortisol levels and inflammatory cytokines help restore physiological stability compromised by chronic stress or cancer-related fatigue [22,23]. These biochemical alterations correspond with improved emotional stability, cognitive clarity, and sleep patterns, which are vital for resilience in individuals facing cancer [24].

Moreover, aerobic activity has been linked to enhanced functional connectivity between the prefrontal cortex and limbic regions in the brain, which supports better top-down emotional control a goal shared by therapeutic practices in psycho-oncology aimed at helping patients manage anxiety and fear through awareness and reappraisal [25]. Furthermore, engagement in aerobic exercise acts as a nonverbal form of cognitive-emotional training, where the physiological adaptations foster self-regulation capabilities [19]. Over time, repeated physical engagement can encourage adaptive stress responses, facilitating a shift from reactive patterns to adaptive coping strategies [18].

Consideration of the autonomic nervous system (ANS) reveals that aerobic exercise not only enhances parasympathetic tone and heart rate variability (HRV) indicators of adaptive flexibility but also positively influences emotional resilience and social engagement [17]. The coordination of breath and movement experienced during aerobic activities cultivates psychophysiological coherence, which is both measurable and teachable, making it an ideal focal point for resilience training within educational frameworks [26].

From a psychological standpoint, exercise contributes significantly to self-efficacy a core concept in resilience theory. Engaging in structured exercise tasks solidifies personal beliefs in one's competence and control, effectively countering feelings of helplessness often experienced during cancer diagnoses and treatments [16]. The progressive nature of aerobic training characterized by goal setting and incremental challenges reflects the psychological processes of adaptation, enabling individuals to confront and integrate challenges rather than evade them [27]. As participants push their physical limits, they glean essential lessons about recovery and personal transformation.

2.3. Integrating movement into educational resilience frameworks

Incorporating aerobic movement into psycho-oncological education provides a real-time embodiment of the biopsychosocial paradigm, allowing various learners patients, caregivers, and healthcare trainees to observe and experience the interactions between movement and emotional states [20]. The educational value stems from embodied cognition, where learning transcends intellectual understanding to incorporate sensory experiences, creating rich, kinesthetic learning opportunities [25]. Through techniques like controlled breathing during physical exertion or interpreting muscle fatigue as a metaphor for emotional endurance, participants engage in experiences that unify physiology with psychological insights [28].

This integrative pedagogical approach resonates with transformative learning theory, which champions reflective, experiential, and embodied learning that fosters meaningful change in the perception of resilience [29]. Aerobic exercise serves as a

safe yet challenging environment where individuals can confront discomfort, reinterpret bodily sensations, and develop narratives of control that empower personal growth [17]. Additionally, in the context of psycho-oncology, where patients often feel alienated from their bodies, reclaiming physical agency through movement can be profoundly therapeutic [24].

Moreover, the integration of movement within psycho-oncological curricula promotes collaborative learning opportunities among professionals, including oncologists, psychologists, physiotherapists, and educators. These collaborations lead to the creation of multidisciplinary modules where physical activities are complemented by reflective dialogues or psychoeducational content, enhancing the overall learning experience [26]. In hybrid and digital formats, utilizing wearable sensors may offer immediate feedback that connects physiological responses to emotional states, thereby deepening understanding and engagement.

Additionally, movement-based education aligns with established mindfulness practices and cognitive-behavioral frameworks within psycho-oncology. While mindfulness typically cultivates awareness through stillness and reflection, aerobic exercise achieves similar outcomes by promoting active engagement, focus, and acceptance in dynamic environments, thereby enhancing pedagogical diversity and addressing varied learning preferences in resiliency training [17].

2.4. Synthesis

Combining the insights from the biopsychosocial and neuropsychobiological models reveals that aerobic exercise encapsulates resilience in practice. It weaves together physiological regulation, psychological flexibility, and social interconnectedness into an experiential learning process [15]. This embodiment of resilience operationalizes it as a skill that can be translated into educational settings. Through movement, learners can authentically experience balance, challenge, and recovery while reflecting on the intertwined nature of their emotional and cognitive processes [16].

By cultivating psycho-oncological education around the lens of movement, the narrative shifts from passive reception to active involvement in personal regulation and resilience-building practices. The following section will illuminate the educational implications of this integration, highlighting how aerobic exercise can serve as a structured pedagogical tool for fostering mind-body resilience in patients, students, and healthcare providers alike.

3. Educational perspectives on movement and resilience

3.1. Psycho-educational models in cancer care

Education in psycho-oncology transcends traditional medical knowledge transfer; it aims to develop self-awareness, coping strategies, and emotional literacy in both patients and healthcare providers. While traditional psycho-educational interventions consisting of lectures, counseling, or cognitive-behavioral sessions have been effective at improving treatment adherence and reducing psychological distress, they often lack a physical component that enables deeper engagement [30]. By

integrating movement-based learning into psycho-education, concepts such as self-efficacy, emotional regulation, and resilience gain a tangible grounding in physical experiences [31].

Cancer can significantly disrupt a patient's sense of bodily autonomy and control. Psycho-educational efforts are designed to restore a sense of coherence between mind and body by equipping patients with the knowledge and tools needed to manage not only painful physical symptoms but also emotional challenges [32]. Aerobic exercise can enhance these objectives by transforming the body from a mere recipient of treatment to an active participant in the healing process. Observing physiological adaptations such as heart rate recovery and increased stamina reinforces the belief that healing emerges from intentional and gradual effort, thus solidifying cognitive beliefs in resilience [33,34].

For students and professionals in healthcare, engaging in structured movement activities fosters empathy by enabling them to vicariously experience fatigue, effort, and self-regulation. These embodied encounters provide a deeper understanding of the psychological principles involved in cancer care, far more effectively than didactic instruction alone [35]. Thus, movement operates as both therapeutic and educational a vital medium for realizing complex psychological constructs through embodied experiences.

3.2. Exercise as pedagogy: Teaching through movement

The educational application of aerobic exercise is grounded in experiential learning theory articulated by Kolb in 1984, which asserts that learning is founded on the transformation of experiences [36]. The cyclical nature of this theory can be directly mapped onto movement practice:

- **Concrete experience:** Participation in aerobic activities like walking, cycling, or group rhythm exercises.
- **Reflective observation:** Increased awareness of body sensations, fatigue levels, and emotional responses.
- **Abstract conceptualization:** Relating these experiences to psychological constructs such as motivation or stress management.
- **Active experimentation:** Adjusting pace, breathing techniques, and mental focus during subsequent sessions.

This model aligns well with the development of psychological resilience, enabling individuals to engage with stress, reflect on their reactions, integrate newly acquired insights, and adapt their behaviors accordingly [37]. Consequently, teaching resilience through movement becomes a dynamic, experiential learning cycle rather than a straightforward transmission of information.

As illustrated in **Figure 1**, the Mind–Body Learning Cycle encompasses four interconnected phases: Movement Experience, Reflection & Awareness, Conceptual Insight, and Application & Adaptation. This process not only supports the understanding of resilience as a skill but also the embodiment of that skill through physical action.



Figure 1. The mind–body learning cycle in psycho-oncological education.

Furthermore, group movement fosters collaborative learning environments, reflecting communal coping mechanisms vital for resilience. Participants learn essential interpersonal skills such as cooperation, shared rhythm, and mutual encouragement, which translate to psychosocial dimensions critical for surviving and thriving through cancer treatment [38].

3.3. Curriculum design for psycho-oncological training

The seamless integration of movement into psycho-oncological education can be adapted across various contexts, including patient survivorship programs, undergraduate medical curricula, and ongoing professional development initiatives [32]. **Table 1** outlines a conceptual model for incorporating aerobic movement into psycho-oncological training. A modular structure integrating aerobic movement into psycho-oncological curricula across patient, professional, and community contexts. The table outlines learning objectives, sample activities, expected psychophysiological outcomes, and assessment methods.

Table 1. Conceptual framework for movement-based psycho-oncological education.

Educational level	Learning objectives	Example activities	Expected psychophysiological outcomes	Assessment methods
Patient/self-management	Develop self-efficacy, body awareness, and emotional regulation	Low-intensity walking groups, breathing-paced cycling	Improved HRV, mood stabilization, increased recovery confidence	Self-reported scales, HRV biofeedback
Medical & nursing education	Cultivate empathy and understanding of mind-body dynamics	Reflective movement sessions integrated into psycho-oncology modules	Enhanced emotional intelligence, improved stress tolerance	Reflective journals, peer discussions
Professional resilience training	Combat burnout and foster mindfulness through aerobic awareness	Group jogging or rhythmic exercise paired with guided reflections	Reduced stress biomarkers, elevated focus	Pre/post psychological well-being scales
Community/public health outreach	Enhance psycho-oncological literacy through participatory education	Community fun-runs, dance-aerobics combined with psycho-education booths	Heightened social cohesion and awareness of psychosocial care	Participation metrics, qualitative feedback

This table exemplifies how movement can be tailored to fit educational contexts while retaining its core intent. Whether as structured classroom exercises or integrated digital modules, the focus remains on experiential and reflective learning.

Digital innovations further enhance accessibility and tracking. Wearable sensors and mobile applications offer real-time biofeedback, linking physiological metrics like heart rate to subjective emotional states. Such tools enable learners to visualize the relationship between physical exertion and emotional regulation, reinforcing academic concepts through data-driven insights [39]. Virtual platforms can also support inclusive education for patients who may be homebound or immunocompromised, ensuring broader access to psycho-oncological training [40].

3.4. Integrative learning environments

Implementing movement-based pedagogy requires collaboration among psychologists, oncologists, physiotherapists, and educators to effectively translate movement into psycho-emotional language. Sessions can begin with brief guided reflection (“Notice your breathing; how does it reflect your emotional state?”), followed by aerobic activities, and conclude with discussions that tie physical sensations to coping principles [41]. Over time, learners can craft personalized narratives of resilience, such as “When my pace steadied, my mind cleared,” or “Fatigue taught me to slow down and breathe,” which can then be articulated through reflective writing or group dialogue, transitioning bodily insights into conceptual understanding [42].

Incorporating art-based or mindfulness exercises like journaling, drawing, or exploring biofeedback results can further integrate cognitive, emotional, and sensory

dimensions into learning [43]. The learning environment should reflect the therapeutic process itself: safe, participatory, and self-directed. The symbolism of rhythm and movement possesses profound psychological implications, as rhythmic synchronization facilitates social bonding and shared emotional regulation. For survivors, moving together embodies solidarity against adversity, while healthcare trainees develop relational empathy through richly embodied experiences [44].

3.5. Challenges and considerations

Despite its potential, integrating aerobic exercise into educational programs poses logistical and theoretical challenges. Prioritizing physical safety, monitoring fatigue levels, and ensuring medical clearance is essential for patient participants. In academic settings, time constraints and institutional inertia could obstruct the incorporation of movement-based education. Furthermore, educators may need to enhance their skills in movement facilitation and interdisciplinary communication to ensure psychological objectives are not lost amid fitness metrics [45].

Cultural considerations are also crucial. Attitudes towards body image, gender roles, and group activities differ widely across cultural contexts. Programs must adapt to these considerations, offering culturally relevant movement forms such as Tai Chi or low-impact dance for those who may resist traditional exercise terminology. The focus remains on ensuring accessibility to the educational intent, aiming to foster resilience and mind-body integration across diverse demographics [46].

Finally, empirical research in this area remains sparse. Most studies to date evaluate exercise outcomes regarding fatigue or quality of life instead of learning outcomes such as self-awareness or emotional competence [45]. Future research should use mixed-method approaches to address both physiological and educational impacts, combining psychometric assessments with qualitative reflections. Establishing such evidence would validate movement-based pedagogy as an innovative educational tool within the realm of psycho-oncology.

3.6. Summary

Educational strategies that embed aerobic movement into psycho-oncology convert this cognitive domain into a practice of embodiment. By aligning physical effort with reflective learning, participants can experience resilience as both a biological rhythm and teachable skill. Whether assisting patients in reclaiming bodily confidence or helping students nurture their empathy, teaching through movement reconnects the human body with its intrinsic adaptability and meaning-making capabilities. The forthcoming chapter will delve into the empirical evidence supporting the psychological and physiological effects of aerobic exercise within cancer populations, highlighting how these mechanisms validate its role as both an educational and therapeutic resource in psycho-oncological care.

4. Evidence linking aerobic exercise to psychological outcomes in cancer

4.1. Overview

A substantial body of evidence indicates that aerobic exercise offers multifaceted benefits for individuals with cancer, extending beyond just cardiorespiratory improvements to include psychological well-being, cognitive clarity, emotional regulation, and biochemical indicators of stress adaptation [47]. Within the context of psycho-oncology, these findings provide a physiological basis for mind-body resilience education. When patients or learners observe measurable changes in physical health alongside improvements in emotional states or coping abilities, it reinforces the understanding that resilience can indeed be cultivated through movement [2,48].

Meta-analyses across various cancer populations including breast, colorectal, hematological, and gynecological cancers demonstrate moderate to large effect sizes for the benefits of exercise on reducing fatigue, anxiety, and depressive symptoms. Structured aerobic programs of moderate intensity (e.g., 30- to 45-minute sessions, 3-5 times per week) have been shown to decrease perceived stress and significantly enhance quality of life compared to usual care [49]. Importantly, improvement correlates with adherence levels and perceived self-efficacy, indicating that how individuals interpret their engagement in exercise can mediate these physiological benefits [34].

4.2. Effects on mood, anxiety, and depression

Cancer-related distress often results from a combination of biological factors, treatment side effects, and existential concerns. Aerobic exercise has been shown to modulate these underlying processes by regulating the hypothalamic-pituitary-adrenal (HPA) axis and promoting the release of endorphins and monoamines [50]. Clinical trials have documented that consistent aerobic activity spanning 8-12 weeks leads to measurable reductions in anxiety and depression, sometimes paralleling the effects of low-dose pharmacological interventions for specific subgroups of patients [51].

Behaviorally, exercise enhances mood regulation through mechanisms such as behavioral activation and the mastery of challenges. Successful completion of exercise sessions offers micro-exposures to stress followed by recovery, reinforcing the belief that emotional states can be influenced positively through purposeful action [52]. Group-based exercise sessions further strengthen these effects by facilitating social connection, which serves as a buffer against depression in cancer patients [53]. In psycho-educational settings, these outcomes can be used to concretely demonstrate the relationship between active participation and emotional control, allowing educators to effectively translate research findings into meaningful pedagogical practices [47].

4.3. Cognitive and neurobehavioral benefits

Chemotherapy, radiation therapy, and chronic stressors can lead to cognitive deficits often referred to as “chemo-brain,” characterized by impaired attention, processing speed, and memory [47]. Aerobic exercise is emerging as a viable, non-

pharmacological intervention against these cognitive impairments. Research utilizing functional MRI has shown that engaging in moderate-intensity aerobic activities can lead to increased hippocampal volume and greater activation in prefrontal cortical regions, which are crucial for executive function and attentional control [34].

From a neuroeducational perspective, these findings signify that participant may authentically experience improved cognitive functions alongside physical activity, exemplifying the embodied cognition concept: as the body undergoes rhythmic movement, the brain undergoes beneficial changes. This dual benefit not only reinforces the physiological grounding of psycho-oncological training but also motivates ongoing participation through direct observation of cognitive improvements [54].

4.4. Physiological correlates of psychological resilience

Regular aerobic exercise has been linked to significant physiological changes, including lower baseline cortisol levels, enhanced parasympathetic tone, and reduced levels of inflammatory cytokines like IL-6 and TNF- α [47]. Improved heart rate variability (HRV), which serves as an indicator of autonomic balance, is consistently associated with positive emotional regulation outcomes and greater resilience to stress among cancer survivors [34].

These physiological markers can offer concrete data for educational biofeedback systems, allowing learners to engage in self-monitoring and regulation practices. For instance, when individuals visualize their HRV improvements in correlation with controlled breathing or moderate exercise, they illustrate resilience as a tangible physiological phenomenon. Such experiences can make abstract psychological constructs more observable and teachable, thereby reinforcing the pedagogical importance of movement [50].

4.5. Social and existential dimensions

Aerobic exercise does not only impact individual health; it also cultivates social connections and promotes existential coherence, both of which are vital components in psycho-oncology. Group exercise fosters a sense of belonging and mutual support, mirroring the communal resilience often found in support groups for cancer patients [48]. As participants engage with one another, they often report increased trust in their bodies and a deeper appreciation for life, correlating with concepts of post-traumatic growth [50,53].

Such collaborative experiences have significant implications for educational practices, illustrating the importance of community resilience where movement is framed as a collective journey rather than a solitary task. In training settings, participation in these group scenarios can enhance empathy and appreciation for the recovery process, allowing healthcare learners to gain deeper insights into their patients' experiences [47].

4.6. Dose–response and practical considerations

Evidence suggests that while moderate-intensity aerobic exercise appears to yield the most significant benefits, the optimal “dose” can vary significantly based on

individual factors such as diagnosis, treatment stage, and fatigue levels. A minimum threshold of approximately 150 minutes of moderate activity per week is associated with positive psychological outcomes, though even shorter bouts of activity (around 10-15 minutes daily) can offer substantial mood benefits [2,51].

Thus, educational programs should focus on promoting individualized approaches to exercise rather than inflexible prescriptions. Learners can be encouraged to monitor their perceived exertion, mood states, and physiological feedback, transforming each exercise session into a personal experiment in self-regulation and emotional monitoring [34,52].

4.7. Integrative summary

In summary, the evidence firmly establishes that aerobic exercise contributes significantly to well-being via multiple interrelated pathways, including biochemical regulation, neurocognitive benefits, behavioral activation, and social engagement. Collectively, these mechanisms form the biological and psychological foundation of resilience. By incorporating these insights into educational frameworks, a comprehensive model emerges for demonstrating how movement influences both physiology and emotional well-being [53].

Figure 2 depicting a layered model physiological mechanisms (e.g., endorphin release, cortisol regulation, HRV improvement) leading to neurocognitive enhancement and psychosocial outcomes such as mood stabilization, self-efficacy, and social connection.




Figure 2. Mechanistic pathways linking aerobic exercise to psychological resilience in cancer.

4.8. Limitations and research gaps

Despite the compelling evidence linking exercise to improved psychosocial outcomes, several limitations remain. Numerous studies prioritize clinical outcomes while neglecting qualitative or educational assessments. Few investigations significantly explore how participants perceive their exercise experiences or how reflective practices might mediate long-term resilience [47,48]. Furthermore, the variability among exercise protocols, adherence levels, and instructional methodologies complicates inter-study comparisons [2].

Future research should adopt hybrid methodologies that merge physiological monitoring with qualitative research to unpack the lived experiences associated with movement in a psycho-oncological context. Investigating how different educational approaches like guided reflections versus unsupervised training affect outcomes could elucidate the pedagogical mechanisms facilitating physiological change [55].

4.9. Conclusion of Chapter 4

The empirical evidence solidifies aerobic exercise as a vital biopsychosocial intervention contributing to improved emotional stability, cognitive functioning, and physiological resilience in cancer patients and survivors. The multidimensional benefits of exercise validate its implementation as not only a therapeutic modality but also a powerful educational tool for fostering mind-body resilience.

In the sphere of psycho-oncology, teaching through movement transforms complex scientific concepts into relatable, felt experiences, bridging molecular changes with meaningful emotional responses. The next chapter will extend this exploration to educational program models and strategies for implementation, aiming to operationalize such evidence in clinical and educational contexts.

5. Educational program models and implementation

5.1. Translating evidence into educational design

Prior chapters have established that aerobic exercise can yield substantial psychophysiological benefits, and these effects can be framed as valuable learning opportunities to foster resilience. This chapter bridges theory and practice by outlining how movement-based psycho-oncological education can be effectively implemented in patient, professional, and community contexts. Successful implementation necessitates an approach that harmonizes clinical safety with educational intent; each movement session operates as both therapeutic and instructional, encouraging participants to recognize the correlation between physiological regulation and emotional recovery [56].

In terms of educational design, this translation follows three progressive stages:

1. **Knowledge acquisition:** Participants engage with the foundational science of exercise and resilience (psycho-education).
2. **Embodied experience:** Learners participate in aerobic movement and observe their physiological and emotional responses.
3. **Reflective integration:** Participants connect physical sensations with psychological concepts through discussions or journaling.

This tri-layer sequence—**Learn** → **Move** → **Reflect**—is adaptable for both clinical and academic environments [57].

5.2. Curriculum architecture for psycho-oncological education

To create a practical curriculum, it is essential to balance theoretical instruction, supervised movement, and reflective learning. **Table 2** proposes a modular framework suitable for adaptation within hospitals, universities, or survivorship centers. The framework follows a stepwise educational design that emphasizes conceptual understanding, guided aerobic practice, reflective dialogue, and continuous feedback each component working together to cultivate mind–body resilience competencies.

Table 2. Modular curriculum framework for teaching mind–body resilience through aerobic exercise.

Module	Focus	Core activities	Duration	Learning outcomes
Foundations of psycho-oncology	Understanding psychosocial stress and resilience	Mini-lectures, case discussions	2 h	Define resilience and explain the biopsychosocial model
Physiology of movement	Exercise science basics for cancer populations	Demonstrations, safety screening	2 h	Recognize physiological limits and safety indicators
Movement practice	Guided aerobic session (walking, cycling, rhythmic flow)	30–40 min	Experience HRV change and mood shift	
Reflection and dialogue	Integrating bodily experience with emotional awareness	Group debrief, reflective journaling	1 h	Translate sensations into coping concepts
Digital feedback & self-monitoring	Using wearables and apps for resilience tracking	Data visualization, HRV/breath tracking	Ongoing	Interpret biofeedback as a learning tool
Evaluation & adaptation	Assessing personal growth and program refinement	Surveys, interviews	End of cycle	Demonstrate improved self-efficacy and emotional regulation

Each module functions independently or can be part of a longer course in psycho-oncology education. For patient programs, shorter cycles (4–6 weeks) with low-intensity sessions are optimal; for students and professionals, extended modules integrating theoretical knowledge and supervision are beneficial [56].

5.3. Interdisciplinary collaboration and instructor roles

Successful implementation relies on interdisciplinary partnerships that bridge oncology, psychology, physiotherapy, and educational theory.

- **Clinical facilitators** are responsible for ensuring exercise safety, monitoring patient fatigue, and modifying intensity according to individual medical conditions.
- **Psychologists or psycho-oncologists** lead reflective and cognitive activities, linking bodily experiences to emotional processes [58].
- **Educators** are tasked with structuring learning objectives, creating assessment rubrics, and designing reflective prompts.

- **Technology specialists** integrate biofeedback tools, ensuring usability and protecting patient data privacy.

Regular team meetings facilitate collaboration and alignment between clinical objectives (e.g., reducing fatigue) and educational outcomes (e.g., enhancing emotional awareness) [59].

5.4. Pedagogical methods and delivery modes

a. Experiential and reflective learning

Movement-based education thrives on the principles of experiential reflection. Participants should document immediate post-activity observations heart rate recovery, breathing patterns, emotional tone and later analyze how these observations correlate with stress regulation [60] Structured prompts like “What did my body teach me today?” or “When did effort feel easy?” can enhance meaning-making and contextualize insights related to mind-body connections.

b. Blended and digital platforms

Hybrid learning environments extend the reach of educational offerings. Online modules can deliver psycho-educational content and self-monitoring tutorials, while in-person sessions provide opportunities for supervised movement and group dialogue. Wearables and smartphone applications can generate quantitative data for example, HRV graphs or mood trackers that reinforce self-efficacy and visible progress [61].

c. Peer-learning models

Peer or survivor mentors may co-facilitate sessions, translating personal experiences into learning opportunities. This shared learning approach fosters authenticity and community cohesion, reflecting the social aspects vital to building resilience [62].

5.5. Program evaluation and quality indicators

Assessing educational success should encompass both clinical and educational metrics.

- **Clinical indicators** may include reductions in fatigue, improved HRV, better sleep quality, and decreased anxiety scores.
- **Educational indicators** may assess increases in self-efficacy, emotional literacy, reflective depth, and knowledge retention.

A mixed-methods assessment strategy combining standardized scales (e.g., Brief Resilience Scale, Hospital Anxiety and Depression Scale) with qualitative feedback provides a holistic view of program effectiveness. Regular feedback loops enable instructors to adjust content’s intensity and emotional weight [63].

5.6. Implementation settings

Clinical rehabilitation units

Hospitals can incorporate 20-minute aerobic education sessions into physiotherapy schedules. A brief pre-exercise orientation can connect movement with coping strategies (e.g., pacing relating to emotional regulation). After each session, participants can complete reflection sheets to capture their insights [64].

University and training programs

In medical and nursing schools, modules can be integrated within psycho-oncology or behavioral science curricula. Students can partake in group sessions that merge physiological insights with psychological concepts, followed by academic reflections that connect bodily feedback to patient empathy and self-care [65].

Community and survivorship programs

Public health centers and survivor networks may host low-intensity walking clubs or rhythmic sessions combined with psycho-educational discussions. These programs shift the focus from rehabilitation to maintaining well-being and fostering community resilience. They also function as outreach initiatives, normalizing conversations around mental health through accessible physical activities [66].

5.7. Barriers and facilitators

Barriers

- Limited availability of time in treatment or academic schedules.
- Insufficient staff training in both facilitative movement techniques and psycho-educational methods.
- Concerns regarding medical safety during active treatment.
- Inconsistent funding or lack of institutional backing.

Facilitators

- Interdepartmental partnerships between rehabilitation and psychology.
- Demonstrated empirical outcomes evidencing quality-of-life improvements attributable to such programs [67].
- Integration of digital monitoring tools to enhance safety and feedback mechanisms.
- Institutional recognition and incentives for educational innovations.

Addressing these barriers requires policy alignment; for instance, medical institutions could classify movement-education sessions as part of psycho-oncological care, or universities might recognize them for academic credit, promoting the legitimacy of “learning through movement” [68].

5.8. Ethical and cultural considerations

Implementation must respect patient autonomy, cultural contexts regarding body dynamics, and various comfort levels concerning group movement activities. Informed consent processes should make it clear that participation is voluntary and that engagement in reflective practices may elicit emotional responses. Additionally, culturally adaptive programs such as Tai Chi or low-impact dance should be offered to maintain inclusivity and support the educational core of self-observation and regulation [69].

5.9. Summary

Introducing aerobic exercise as a teaching method in psycho-oncology transforms conventional rehabilitation into a journey of resilience cultivation. Through thoughtfully structured curricula, interdisciplinary delivery, and integrated reflective practices, participants not only enhance their physical capabilities but also develop

psychological literacy and adaptive skills. Educational initiatives grounded in movement encapsulate the biopsychosocial model: knowledge transforms into experience, emotions are regulated, and the body serves both as classroom and instructor. Future practice must continually refine these frameworks through systematic evaluations, ensuring that teaching mind-body resilience through movement remains a dynamic and evidence-supported component of psycho-oncological education.

6. Challenges, gaps, and future directions

6.1. Conceptual and methodological gaps

Although the therapeutic and educational promise of aerobic exercise in psycho-oncology is compelling, its integration into formal educational frameworks remains conceptually and methodologically underdeveloped. Most existing studies approach exercise as a clinical intervention, emphasizing outcomes such as fatigue reduction, increase in Quality of Life (QoL) or cardiovascular improvement [70,71], while few investigate exercise as a pedagogical process through which resilience and self-regulation are consciously learned.

This conceptual gap limits understanding of *how* movement generates psychological meaning. Without clear educational models, exercise risks being reduced to a prescriptive health behavior rather than a transformative learning medium. Future frameworks should therefore bridge exercise science with learning theory linking physiological adaptation to cognitive and reflective outcomes.

Methodologically, heterogeneity in study design also constrains evidence synthesis. Differences in cancer type, treatment stage, and supervision create inconsistent findings. Most interventions employ quantitative measures such as heart-rate variability (HRV) [72,73] or depression scales [74–76], yet seldom integrate qualitative reflections that capture participants' lived experiences of movement. Recent work using HRV monitoring demonstrates promising links between autonomic regulation and resilience outcomes [77,78], but the educational translation of these biological signals remains limited. Narrative and mixed-methods approaches could reveal the *inner learning process* underlying these physiological adaptations.

6.2. Implementation barriers in educational contexts

Despite growing endorsement of holistic cancer care, educational integration faces multiple barriers.

1. **Institutional inertia**—oncology and academic programs often separate physical rehabilitation from psychosocial education.
2. **Resource and training constraints**—implementing movement-based education requires instructors versed in both physiology and psycho-education, a rare combination.
3. **Safety and liability concerns**—medical oversight is essential, particularly during treatment.
4. **Time and space limitations**—embodied learning sessions compete with dense clinical and curricular schedules.

5. **Evaluation challenges**—emotional literacy and self-efficacy are difficult to quantify using conventional biomedical metrics.

Addressing these issues requires systemic redesign and interdisciplinary collaboration between educators, physiotherapists, and psycho-oncologists. Standardized modules, supported by empirical data on exercise safety and efficacy, can help institutionalize the approach.

6.3. Cultural and ethical considerations

Cultural norms strongly influence attitudes toward exercise, body image, and group participation. In some regions, exertion during illness may be stigmatized or gender-restricted. Hence, culturally adaptive practices such as Tai Chi, rhythmic walking, or low-impact dance may convey the same educational principles while aligning with community identity.

Ethically, programs must prioritize informed consent and emotional safety. Because movement can evoke vulnerability or body memories, facilitators should be trained to respond with empathy and confidentiality. Reflection activities must remain voluntary, ensuring psychological boundaries are respected.

6.4. Research gaps and opportunities

Several key research directions emerge from current literature:

- **Longitudinal learning retention.**
 - Most trials measure short-term benefits; few assess whether participants maintain resilience skills over time.
- **Comparative pedagogy.**
 - Research comparing lecture-based psycho-education with embodied, movement-based learning could determine which better enhances emotional and cognitive outcomes.
- **Integration of neurobiological and reflective data.**
 - Combining biomarkers (e.g., HRV, BDNF) with reflective journals would demonstrate how awareness mediates physiological change.
- **Technology-assisted learning analytics.**
 - Wearables and AI analytics can personalize pacing and track adaptation, but studies on data ethics and learner autonomy remain scarce.
- **Instructor development.**
 - Little evidence exists regarding how educators themselves acquire competence in teaching through movement; professional-development research is essential.

6.5. Technological and innovation pathways

Digital transformation now enables scalable and individualized resilience education. Wearable sensors track HRV and stress responses, linking biological feedback with psycho-educational reflection. Mobile applications provide guided movement sessions and emotional logging. Virtual-reality platforms allow patients with mobility limitations to rehearse movement and regulate breathing in immersive environments.

However, technology must serve pedagogy not dominate it. The goal is to enhance self-awareness rather than outsource regulation to devices. Research integrating digital monitoring with reflection may reveal how data visualization strengthens motivation and understanding [77]. Ethical governance is vital to ensure privacy and prevent over-medicalization of learning.

6.6. Policy and institutional directions

Embedding movement-based psycho-oncological education within healthcare systems requires policy recognition:

- **Academic policy:** integrate experiential resilience modules into health-science curricula.
- **Clinical policy:** recognize movement education as reimbursable psycho-social care.
- **Public-health policy:** promote community “mind–body literacy” campaigns linking exercise and psychological well-being.

Collaborative partnerships among hospitals, universities, and NGOs can establish accreditation for movement educators, ensuring consistency and patient safety.

6.7. Future conceptual evolution: toward an “embodied curriculum”

The long-term vision extends toward an embodied curriculum in psycho-oncology. In such a model, movement itself becomes a teaching language. Lessons on stress, emotion, and coping are conveyed through physical experiences that mirror psychological adaptation.

This paradigm redefines both teaching and healing. Educators act as facilitators of body–mind dialogue, and learners become active participants in their own resilience. Conceptually, it operationalizes the biopsychosocial model into lived practice. Initiatives such as *Mind–Body Resilience Labs* could unite research, training, and community outreach, transforming psycho-oncology education into an ongoing, interactive process of physiological and emotional learning.

6.8. Summary and forward vision

The intersection of psycho-oncology, movement science, and education heralds a new era of holistic care. The remaining gaps methodological, institutional, and cultural highlight the need for longitudinal, interdisciplinary, and ethically guided innovation.

Future work should shift from asking *whether* exercise improves mental health to exploring *how learners internalize resilience through movement*. This transition situates aerobic education within the continuum of healing, knowledge, and human development.

Ultimately, the next frontier is not to prescribe exercise as treatment but to teach through movement turning each breath, rhythm, and heartbeat into a living lesson in adaptation, connection, and hope [79].

7. Conceptual framework and conclusion

7.1. Integrative overview

The preceding chapters have traced a continuum from theory to practice: beginning with the biopsychosocial foundations of psycho-oncology, extending through the neuropsychobiological effects of exercise, and culminating in educational applications that teach resilience through embodied learning.

Together, these strands converge into a new synthesis the Educational Pathway of Mind–Body Resilience, which positions aerobic movement not merely as a therapeutic adjunct, but as a *pedagogical process* that unites physiological adaptation, psychological growth, and educational empowerment.

In this integrated view, movement becomes a medium for learning self-regulation. Each exercise session functions as a microcosm of coping a lived rehearsal of stress, effort, and recovery. When participants reflect on these experiences, they transform physical feedback into emotional insight and cognitive understanding. This iterative cycle, in which experience informs awareness and awareness shapes adaptation, lies at the heart of the framework.

7.2. The educational pathway of mind–body resilience

Figure 3 presents the conceptual framework illustrating how aerobic movement functions as an experiential educational pathway for developing mind–body resilience within psycho-oncology. The model comprises four progressive and interlinked stages: Physiological Activation, Cognitive–Emotional Awareness, Reflective Integration, and Educational Transformation positioned vertically to signify the gradual deepening of learning from bodily experience to psychological understanding.




Figure 3. ConceptTUAL FRAmework: The educational pathway of mind–body resilience.

- **Physiological Activation** represents the biological entry point where rhythmic aerobic movement regulates stress systems through endorphin release, heart-rate variability improvement, and neurocognitive stimulation.
- **Cognitive–Emotional Awareness** emerges as participants recognize how bodily rhythms influence mood and thought, building insight into the mind–body interface.
- **Reflective Integration** transforms awareness into meaning via guided reflection, journaling, or dialogue, allowing learners to connect physiological sensations with resilience principles such as acceptance and control.
- **Educational Transformation** marks the point where embodied insights become transferable knowledge, fostering self-efficacy, empathy, and adaptive coping in both patients and healthcare learners.

Two vertical pillars, *Instructor Facilitation* and *Social Context*, flank the process. They emphasize the educator’s role in guiding safe reflection and the group’s role in reinforcing motivation, belonging, and shared resilience. The caption below the figure summarizes the overall premise:

“Aerobic movement as an experiential curriculum linking body regulation, emotional literacy, and resilience learning.”

Together, these elements portray aerobic exercise not as passive rehabilitation but as an active educational medium, demonstrating how mind–body synchronization, reflection, and social interaction collectively nurture resilience in psycho-oncological education

Stage 1: Physiological activation

This stage represents the biological entry point of learning. Through rhythmic aerobic activity, the cardiovascular and neuroendocrine systems engage in stress regulation releasing endorphins, balancing cortisol, and increasing cerebral oxygenation. These responses constitute the *embodied data* from which learners later derive meaning. Physiological coherence (e.g., heart-rate variability, regulated breathing) creates the internal stability necessary for emotional processing.

Empirical studies affirm that such responses are measurable and reproducible. Increases in HRV and decreases in inflammatory markers following structured exercise [77,78] illustrate how biological modulation supports resilience. Within education, this stage sets the groundwork for learning by feeling perceiving one’s own body as a responsive system rather than a passive vessel.

Stage 2: Cognitive–emotional awareness

As physiological regulation stabilizes, attention shifts toward awareness *recognizing* the link between body and mind. Participants begin to observe sensations, thoughts, and emotional shifts during exercise. This metacognitive recognition mirrors therapeutic mindfulness, but it occurs in motion. Educators facilitate this by prompting reflection-in-action:

“Notice when your breath steadies what changes in your thoughts?”

Awareness transforms raw sensory data into conscious understanding. Learners perceive those physiological states influence emotional tone, demonstrating the bidirectional communication central to the biopsychosocial model.

Stage 3: Reflective integration

Reflection converts awareness into personal meaning. After activity, structured dialogues or journaling guide participants to interpret bodily experience through psychological frameworks: endurance as perseverance, rhythm as emotional regulation, fatigue as a cue for rest. This step completes the experiential learning cycle linking concrete experience to abstract insight.

Through guided reflection, individuals internalize resilience principles such as acceptance, pacing, and cognitive flexibility. In educational contexts, reflective debriefings reinforce conceptual learning about stress physiology and coping mechanisms. As shown in community exercise and education programs [80], group reflection also fosters shared learning and empathy, aligning with psycho-oncological principles of communal support.

Stage 4: Educational transformation

At this final stage, embodied insight becomes transferable knowledge applicable to daily life, clinical training, or survivorship. Learners evolve from passive recipients of care to active co-creators of well-being. For patients, transformation manifests as increased agency and self-efficacy; for students and healthcare professionals, it signifies deeper empathy and professional resilience.

Educational transformation also feeds back into institutional culture. When movement-based pedagogy is embedded within oncology curricula or rehabilitation programs, it shifts the paradigm from “treating disease” to “teaching recovery.” This stage exemplifies the educational mission of psycho-oncology: empowering individuals to understand and regulate their internal systems as part of lifelong adaptation.

7.3. Interconnecting mechanisms

The four stages operate within a dynamic system moderated by two contextual pillars:

- **Instructor Facilitation:** Skilled educators act as translators between physiology and psychology, guiding awareness, maintaining safety, and nurturing reflection. Their role mirrors that of a therapist-educator, enabling cognitive and emotional bridging.
- **Social Context:** Group movement amplifies psychosocial outcomes. Shared rhythm enhances belonging, while collective reflection builds empathy and solidarity. Social bonding observed in aerobic community programs confirms the interpersonal dimension of resilience learning [81].

These pillars ensure that learning remains both structured and humane scientifically grounded yet emotionally inclusive.

7.4. Practical implications

The framework offers actionable pathways for curriculum design, research, and policy.

1. **Curricular integration:** Educational institutions can embed short movement-reflection cycles within psycho-oncology training. For instance, weekly 20-

minute guided aerobic sessions followed by 10-minute reflective discussions reinforce psychophysiological principles experientially.

2. **Research methodology:** Future studies should employ mixed-methods designs combining physiological data (HRV, cortisol) with qualitative reflections to validate the four-stage learning process.
3. **Program evaluation:** Success metrics should measure not only clinical outcomes (fatigue, anxiety) but also educational outcomes self-awareness, emotional literacy, and resilience confidence.
4. **Policy development:** Hospitals and universities should institutionalize “movement-education sessions” as recognized psycho-social care components, paralleling models in community survivorship programs [82,83]

7.5. Future research directions

To advance the evidence base, research should address:

- **Mechanistic mapping:** How do neurobiological changes translate into sustained cognitive emotional growth?
- **Pedagogical standardization:** What instructor training models best convey movement-based learning safely?
- **Cultural adaptation:** How can this framework be localized (e.g., integrating walking meditation or Tai Chi for Asian contexts)?
- **Digital embodiment:** Can wearable feedback (HRV apps, VR training) support awareness and reflection without diluting authenticity?

These questions will refine the educational precision and cross-cultural scalability of movement-based psycho-oncology.

8. Conclusion

This narrative review highlights the emerging educational value of aerobic exercise within psycho-oncology, reframing movement as more than a rehabilitative adjunct. Aerobic activity through its physiological, psychological, and social dimensions offers a dynamic platform for teaching mind–body resilience. Evidence consistently shows that structured aerobic programs enhance mood regulation, reduce fatigue, and improve cognitive and emotional well-being among cancer patients and survivors.

By integrating these effects into psycho-oncological education, movement becomes both method and message: learners experience in real time how regulation of the body fosters stability of the mind. The proposed framework of Educational Pathway of Mind–Body Resilience positions aerobic exercise as a cyclical learning process involving physiological activation, emotional awareness, reflection, and educational transformation.

Implementing this model requires interdisciplinary collaboration, culturally sensitive adaptation, and thoughtful instructor facilitation. When embedded in clinical rehabilitation, academic curricula, or community programs, aerobic-based learning can strengthen self-efficacy, empathy, and psychological adaptability across diverse populations.

Ultimately, teaching resilience through movement unites science, education, and healing. It embodies the biopsychosocial philosophy of psycho-oncology demonstrating that recovery and learning are not separate processes but reciprocal acts of growth. As the field continues to evolve, integrating aerobic exercise as an educational practice may redefine how both patients and professionals understand and cultivate resilience: not only to survive illness, but to move through it with awareness, strength, and purpose.

Funding: This research received no external funding.

Ethical approval: Not applicable.

Informed consent statement: Not applicable.

Conflict of interest: The authors declare no conflict of interest.

References

1. Meoded MD, Tănase M, Mehedințu C, Cirimbei C. Psycho-oncology in breast cancer: Supporting women through distress, treatment, and recovery—Three arguments—Rapid narrative review. *Medicina*. 2025; 61(6): 1008. doi: 10.3390/medicina61061008
2. Rosberger Z, Perez S, Bloom J, et al. The missing piece: Cancer prevention within psycho-oncology—A commentary. *Psycho-Oncology*. 2015; 24(10): 1330-1337. doi: 10.1002/pon.3916
3. Ilgen J, Renovanz M, Stengel A, et al. Resilience as a factor influencing psychological distress experience in patients with neuro-oncological disease. *Current Oncology*. 2022; 29(12): 9875-9883. doi: 10.3390/curroncol29120776
4. Ludolph P, Kunzler AM, Stoffers-Winterling J, et al. Interventions to promote resilience in cancer patients. *Deutsches Ärzteblatt International*. 2019; 116: 865-872. doi: 10.3238/arztebl.2019.0865
5. Yu K, Yin B, Zhu Y, et al. Efficacy of a digital postoperative rehabilitation intervention in patients with primary liver cancer: Randomized controlled trial. *JMIR mHealth and uHealth*. 2025; 13: e59228. doi: 10.2196/59228
6. Hall CC, Cook J, Maddocks M, et al. Combined exercise and nutritional rehabilitation in outpatients with incurable cancer: A systematic review. *Supportive Care in Cancer*. 2019; 27(7): 2371-2384. doi: 10.1007/s00520-019-04749-6
7. Qi Y, Wang M, Xue Y, et al. Feasibility of an exercise-nutrition-psychology integrated rehabilitation model based on mobile health and virtual reality for cancer patients: A single-center, single-arm, prospective phase II study. *BMC Palliative Care*. 2024; 23(1): 155. doi: 10.1186/s12904-024-01487-3
8. Gilchrist SC, Barac A, Ades PA, et al. Cardio-oncology rehabilitation to manage cardiovascular outcomes in cancer patients and survivors: A scientific statement from the American Heart Association. *Circulation*. 2019; 139(21). doi: 10.1161/CIR.0000000000000679
9. Bennett AE, O'Neill L, Connolly D, et al. Patient experiences of a physiotherapy-led multidisciplinary rehabilitative intervention after successful treatment for oesophago-gastric cancer. *Supportive Care in Cancer*. 2018; 26(8): 2615-2623. doi: 10.1007/s00520-018-4112-6
10. O'Neill L, Guinan E, Doyle SL, et al. Rehabilitation strategies following esophageal cancer (the ReStOre trial): A feasibility study. *Diseases of the Esophagus*. 2017; 30(5): 1-8. doi: 10.1093/dote/dow012
11. Sabani N, Hardaker G, Sabki A, Salleh S. Understandings of Islamic pedagogy for personalised learning. *The International Journal of Information and Learning Technology*. 2016; 33(2): 78-90. doi: 10.1108/IJILT-01-2016-0003
12. Garrett F. Engaged pedagogy through role-play in a Buddhist studies classroom. *Teaching Theology & Religion*. 2018; 21(4): 336-344. doi: 10.1111/teth.12462
13. Nguyen DJ, Larson JB. Don't forget about the body: Exploring the curricular possibilities of embodied pedagogy. *Innovative Higher Education*. 2015; 40(4): 331-344. doi: 10.1007/s10755-015-9319-6
14. Senf B, Fettel J, Demmerle C, Maiwurm P. Physicians' attitudes towards psycho-oncology, perceived barriers, and psychosocial competencies: Indicators of successful implementation of adjunctive psycho-oncological care? *Psycho-Oncology*. 2019; 28(2): 415-422. doi: 10.1002/pon.4962

15. Brown JC, Huedo-Medina TB, Pescatello LS, et al. The efficacy of exercise in reducing depressive symptoms among cancer survivors: A meta-analysis. *PLoS One*. 2012; 7(1): e30955. doi: 10.1371/journal.pone.0030955
16. Zheng D, Marbut AR, Zhang J, O’Keefe LC. The contribution of psychological resilience and job meaningfulness to well-being of working cancer survivors. *Workplace Health & Safety*. 2022; 70(10): 468-478. doi: 10.1177/21650799221085466
17. Ağaç M, Üzar-Özçetin YS. Psychological resilience, metacognitions, and fear of recurrence among cancer survivors and family caregivers. *Cancer Nursing*. 2022; 45(2): E454-E462. doi: 10.1097/NCC.0000000000000973
18. Fernandes JB, Domingos J, Almeida AS, et al. Enablers, barriers and strategies to build resilience among cancer survivors: A qualitative study protocol. *Frontiers in Psychology*. 2023; 14. doi: 10.3389/fpsyg.2023.1049403
19. Clark V, Brown RF, Gada U, Kamen CS. Resilience related to anxiety, self-esteem, and internalized lesbian, gay, bisexual, and/or transgender negativity: A stepwise regression model among lesbian, gay, bisexual, transgender, queer, and/or another gender and/or sexually expansive identity cancer survivors. *Translational Issues in Psychological Science*. 2025; 11(1): 107-118. doi: 10.1037/tps0000451
20. Park JH, Lee JS, Ko YH, Kim YH. Physical activity of Korean cancer survivors is associated with age and sex. *The Korean Journal of Internal Medicine*. 2021; 36(Suppl 1): S225-S234. doi: 10.3904/kjim.2019.240
21. Javellana M, Hlubocky FJ, Somasegar S, et al. Resilience in the face of pandemic: The impact of COVID-19 on the psychologic morbidity and health-related quality of life among women with ovarian cancer. *JCO Oncology Practice*. 2022; 18(6): e948-e957. doi: 10.1200/OP.21.00514
22. Lim J, Shon E, Paek M, Daly B. The dyadic effects of coping and resilience on psychological distress for cancer survivor couples. *Supportive Care in Cancer*. 2014; 22(12): 3209-3217. doi: 10.1007/s00520-014-2334-9
23. Papathanasiou IV, Kelepouris K, Valari C, et al. Depression, anxiety and stress among patients with hematological malignancies and the association with quality of life: A cross-sectional study. *Medicine and Pharmacy Reports*. 2020; 93(1): 62-68. doi: 10.15386/mpr-1502
24. Üzar-Özçetin YS, Kovanci MS. “For Years I lived in a Cage. Now the Cage Is Open and I Am Just Learning to Fly.” Perspectives of cancer survivors on psychological resilience. *Cancer Nursing*. 2022; 45(4): 297-305. doi: 10.1097/NCC.0000000000001039
25. Kim M, Sok S. Factors influencing resilience among breast cancer survivors: Implications for evidence-based practice. *Worldviews on Evidence-Based Nursing*. 2024; 21(1): 87-95. doi: 10.1111/wvn.12678
26. Tarasenko Y, Chen C, Schoenberg N. Self-reported physical activity levels of older cancer survivors: Results from the 2014 national health interview survey. *Journal of the American Geriatrics Society*. 2017; 65(2): e39-e44. doi: 10.1111/jgs.14589
27. Jones LW, Courneya KS. Exercise counseling and programming preferences of cancer survivors. *Cancer Practice*. 2002; 10(4): 208-215. doi: 10.1046/j.1523-5394.2002.104003.x
28. Sprod LK, Hsieh CC, Hayward R, Schneider CM. Three versus six months of exercise training in breast cancer survivors. *Breast Cancer Research and Treatment*. 2010; 121(2): 413-419. doi: 10.1007/s10549-010-0913-0
29. Smith-Turchyn J, Allen L, Dart J, et al. Characterizing the exercise behaviour, preferences, barriers, and facilitators of cancer survivors in a rural Canadian community: A cross-sectional survey. *Current Oncology*. 2021; 28(4): 3172-3187. doi: 10.3390/currenol28040276
30. Dams L, Van der Gucht E, Devoogdt N, et al. Effect of pain neuroscience education after breast cancer surgery on pain, physical, and emotional functioning: A double-blinded randomized controlled trial (EduCan trial). *Pain*. 2023; 164(7): 1489-1501. doi: 10.1097/j.pain.0000000000002838
31. Wick K, Leeger-Aschmann CS, Monn ND, et al. Interventions to promote fundamental movement skills in childcare and kindergarten: A systematic review and meta-analysis. *Sports Medicine*. 2017; 47(10): 2045-2068. doi: 10.1007/s40279-017-0723-1
32. Ernstmann N, Enders A, Halbach S, et al. Psycho-oncology care in breast cancer centres: A nationwide survey. *BMJ Supportive & Palliative Care*. 2020; 10(4): e36. doi: 10.1136/bmjspcare-2018-001704
33. Lee BO, Choi WJ, Sung NY, et al. Incidence and risk factors for psychiatric comorbidity among people newly diagnosed with cancer based on Korean national registry data. *Psycho-oncology*. 2015; 24(12): 1808-1814. doi: 10.1002/pon.3865
34. Kang JI, Sung NY, Park SJ, et al. The epidemiology of psychiatric disorders among women with breast cancer in South Korea: Analysis of national registry data. *Psycho-oncology*. 2014; 23(1): 35-39. doi: 10.1002/pon.3369
35. Hui D, Kim SH, Kwon JH, et al. Access to palliative care among patients treated at a comprehensive cancer center. *The Oncologist*. 2012; 17(12): 1574-1580. doi: 10.1634/theoncologist.2012-0192

36. Singer S, Dieng S, Wesselmann S. Psycho-oncological care in certified cancer centres—A nationwide analysis in Germany. *Psycho-oncology*. 2013; 22(6): 1435-1437. doi: 10.1002/pon.3145
37. Hammon DE, Pearsall K, Smith NM, et al. Eutectic mixture of lidocaine and prilocaine decreases movement and propofol requirements for pediatric lumbar puncture during deep sedation: A randomized, placebo-controlled, double blind trial. *Journal of Pediatric Hematology/Oncology*. 2022; 44(1): e213-e216. doi: 10.1097/MPH.0000000000002169
38. Zhu H, Chen S, Xie P, et al. Quality of reporting in oncology randomized controlled trials. *Cancer Control*. 2018; 25(1). doi: 10.1177/1073274818781309
39. Fernández-Rodríguez EJ, González-Sánchez J, Puente-González AS, et al. Specific autonomy recovery programme in a comprehensive rehabilitation on functionality and respiratory parameters in oncological patients with dyspnoea. Study protocol. *BMC Nursing*. 2021; 20(1): 120. doi: 10.1186/s12912-021-00633-z
40. Bakitas M, Lyons KD, Hegel MT, et al. Effects of a palliative care intervention on clinical outcomes in patients with advanced cancer: The Project ENABLE II randomized controlled trial. *JAMA*. 2009; 302(7): 741-749. doi: 10.1001/jama.2009.1198
41. Chuck KW, Hoang T, Kim J. Patterns of oncological clinical trials in African countries: A scoping review. *Journal of Cancer Prevention*. 2025; 30(3): 154-165. doi: 10.15430/JCP.25.021
42. McClelland S. Accuracy of medical oncology prognosis for metastatic cancer patients evaluated for enrollment onto an ongoing randomized clinical trial. *American Journal of Clinical Oncology*. 2024; 47(10): 496-497. doi: 10.1097/COC.0000000000001122
43. Windisch P, Dennstädt F, Koechli C, et al. A pipeline for the automatic identification of randomized controlled oncology trials and assignment of tumor entities using natural language processing. *Oncology*. 2024. doi: 10.1101/2024.07.01.24309767
44. Margevicius S, Daly B, Schluchter M, et al. Randomized trial of a web-based nurse education intervention to increase discussion of clinical trials. *Contemporary Clinical Trials Communications*. 2021; 22: 100789. doi: 10.1016/j.conctc.2021.100789
45. Carmona-Gonzalez CA, Cunha MT, Menjak IB. Bridging research gaps in geriatric oncology: Unraveling the potential of pragmatic clinical trials. *Current Opinion in Supportive and Palliative Care*. 2024; 18(1): 3-8. doi: 10.1097/SPC.0000000000000688
46. Olivier T, Haslam A, Prasad V. Reporting of physicians' or investigators' choice of treatment in oncology randomized clinical trials. *JAMA Netw Open*. 2022; 5(1): e2144770. doi: 10.1001/jamanetworkopen.2021.44770
47. Antoni MH. Psychosocial intervention effects on adaptation, disease course and biobehavioral processes in cancer. *Brain, Behavior, and Immunity*. 2013; 30: S88-S98. doi: 10.1016/j.bbi.2012.05.009
48. Jacobsen PB. Clinical practice guidelines for the psychosocial care of cancer survivors: Current status and future prospects. *Cancer*. 2009; 115(S18): 4419-4429. doi: 10.1002/cncr.24589
49. Baziliansky S, Cohen M. Emotion regulation and psychological distress in cancer survivors: A systematic review and meta-analysis. *Stress and Health*. 2021; 37(1): 3-18. doi: 10.1002/smi.2972
50. Namjoshi S. The crucial role of psychosocial research for patients and caregivers: A narrative review of pediatric psycho-oncology research in India. *Psycho-Oncology Research in Pediatric Cancer in India*. 2024; 45(3): 202-209. doi: 10.1055/s-0044-1786967
51. Conley CC, Bishop BT, Andersen BL. Emotions and emotion regulation in breast cancer survivorship. *Healthcare*. 2016; 4(3): 56. doi: 10.3390/healthcare4030056
52. Mock V, St. Ours C, Hall S, et al. Using a conceptual model in nursing research—Mitigating fatigue in cancer patients. *Journal of Advanced Nursing*. 2007; 58(5): 503-512. doi: 10.1111/j.1365-2648.2007.04293.x
53. Turner J. The changing landscape of cancer care—The impact of psychosocial clinical practice guidelines. *Psycho-Oncology*. 2015; 24(4): 365-370. doi: 10.1002/pon.3803
54. Miller M, Vachon E, Kwekkeboom K. Cancer-related symptom frameworks using a biopsychosocial-spiritual perspective: A scoping review. *Western Journal of Nursing Research*. 2023; 45(10): 963-973. doi: 10.1177/01939459231193698
55. Pradeu T, Daignan-Fornier B, Ewald A, et al. Reuniting philosophy and science to advance cancer research. *Biological Reviews*. 2023; 98(5): 1668-1686. doi: 10.1111/brv.12971
56. Aburizik A, Raque TL, Spitz N, et al. Responding to distress in cancer care: Increasing access to psycho oncology services through integrated collaborative care. *Psycho-Oncology*. 2023; 32(11): 1675-1683. doi: 10.1002/pon.6217

57. John JM, Haug V, Thiel A. Physical activity behavior from a transdisciplinary biopsychosocial perspective: A scoping review. *Sports Medicine - Open*. 2020; 6(1): 49. doi: 10.1186/s40798-020-00279-2
58. Meggiolaro E, De Padova S, Ruffilli F, et al. From distress screening to uptake: An Italian multicenter study of cancer patients. *Cancers*. 2021; 13(15): 3761. doi: 10.3390/cancers13153761
59. Dilworth S, Higgins I, Parker V, et al. Patient and health professional's perceived barriers to the delivery of psychosocial care to adults with cancer: A systematic review. *Psycho-oncology*. 2014; 23(6): 601-612. doi: 10.1002/pon.3474
60. Lawson McLean AC, Lawson McLean A, Ernst T, et al. Benchmarking palliative care practices in neurooncology: A German perspective. *Journal of Neuro-Oncology*. 2024; 168(2): 333-343. doi: 10.1007/s11060-024-04674-7
61. Grassi L, Watson M. Psychosocial care in cancer: An overview of psychosocial programmes and national cancer plans of countries within the International Federation of Psycho-Oncology Societies. *Psycho-oncology*. 2012; 21(10): 1027-1033. doi: 10.1002/pon.3154
62. Anuk D, Özkan M, Kizir A, Özkan S. The characteristics and risk factors for common psychiatric disorders in patients with cancer seeking help for mental health. *BMC Psychiatry*. 2019; 19(1): 269. doi: 10.1186/s12888-019-2251-z
63. Brebach R, Sharpe L, Costa DSJ, et al. Psychological intervention targeting distress for cancer patients: A meta-analytic study investigating uptake and adherence. *Psycho-oncology*. 2016; 25(8): 882-890. doi: 10.1002/pon.4099
64. Brédart A, Didier F, Robertson C, et al. Psychological distress in cancer patients attending the European Institute of Oncology in Milan. *Oncology*. 1999; 57(4): 297-302. doi: 10.1159/000012064
65. Tada Y, Matsubara M, Kawada S, et al. Psychiatric disorders in cancer patients at a university hospital in Japan: Descriptive analysis of 765 psychiatric referrals. *Japanese Journal of Clinical Oncology*. 2012; 42(3): 183-188. doi: 10.1093/jjco/hyr200
66. Kadan-Lottick NS, Vanderwerker LC, Block SD, et al. Psychiatric disorders and mental health service use in patients with advanced cancer. *Cancer*. 2005; 104(12): 2872-2881. doi: 10.1002/cncr.21532
67. Mirzaie K, Burns-Gebhart A, Meyerheim M, et al. Emotion regulation flexibility and electronic patient-reported outcomes: A framework for understanding symptoms and affect dynamics in pediatric psycho-oncology. *Cancers*. 2022; 14(16): 3874. doi: 10.3390/cancers14163874
68. Donovan KA, Grassi L, Deshields TL, et al. Advancing the science of distress screening and management in cancer care. *Epidemiology and Psychiatric Sciences*. 2020; 29: e85. doi: 10.1017/S2045796019000799
69. Kusch M, Labouvie H, Schiewer V, et al. Integrated, cross-sectoral psycho-oncology (isPO): A new form of care for newly diagnosed cancer patients in Germany. *BMC Health Services Research*. 2022; 22(1): 543. doi: 10.1186/s12913-022-07782-0
70. Sibarani LP, Setijadi AR, Apriningsih H, et al. Effects of aerobic exercise on the six minutes walking test and quality of life in EGFR mutation non-small cell lung cancer patients. *Jurnal Respirologi Indonesia*. 2025; 45(2): 93-99. doi: 10.36497/jri.v45i2.613
71. Courneya KS, Sellar CM, Stevinson C, et al. Randomized controlled trial of the effects of aerobic exercise on physical functioning and quality of life in lymphoma patients. *Journal of Clinical Oncology*. 2009; 27(27): 4605-4612. doi: 10.1200/JCO.2008.20.0634
72. Niederer D, Vogt L, Thiel C, et al. Exercise effects on HRV in cancer patients. *International Journal of Sports and Medicine*. 2012; 34(01): 68-73. doi: 10.1055/s-0032-1314816
73. Niederer D, Vogt L, Gonzalez-Rivera J, et al. Heart rate recovery and aerobic endurance capacity in cancer survivors: Interdependence and exercise-induced improvements. *Supportive Care in Cancer*. 2015; 23(12): 3513-3520. doi: 10.1007/s00520-015-2719-4
74. Aydin M, Kose E, Odabas I, et al. The effect of exercise on life quality and depression levels of breast cancer patients. *Asian Pacific Journal of Cancer Prevention*. 2021; 22(3): 725-732. doi: 10.31557/APJCP.2021.22.3.725
75. Mehnert A, Veers S, Howaldt D, et al. Effects of a physical exercise rehabilitation group program on anxiety, depression, body image, and health-related quality of life among breast cancer patients. *Onkologie*. 2011; 34(5): 248-253. doi: 10.1159/000327813
76. Kulchyski M, Halder HR, Askin N, et al. Aerobic physical activity and depression among patients with cancer. *JAMA Network Open*. 2024; 7(10): e2437964. doi: 10.1001/jamanetworkopen.2024.37964
77. Grote S, Ricci JM, Dehom S, et al. Heart rate variability and cardiovascular adaptations among cancer-survivors following a 26-week exercise intervention. *Integrative Cancer Therapies*. 2020; 19. doi: 10.1177/1534735420969816
78. Kloter E, Barrueto K, Klein SD, et al. Heart rate variability as a prognostic factor for cancer survival—A systematic review. *Frontiers in Physiology*. 2018; 9. doi: 10.3389/fphys.2018.00623

79. Zhu G, Zhang X, Wang Y, et al. Effects of exercise intervention in breast cancer survivors: A meta-analysis of 33 randomized controlled trials. *OncoTargets and Therapy*. 2016; 2016: 2153-5168. doi: 10.2147/OTT.S97864
80. Musanti R, Murley B. Community-based exercise programs for cancer survivors. *Clinical Journal of Oncology Nursing*. 2016; 20(6): S25-S30. doi: 10.1188/16.CJON.S2.25-30
81. Alberton CL, Andrade LS, Xavier BEB, et al. Land- and water-based aerobic exercise program on health-related outcomes in breast cancer survivors (WaterMama): Study protocol for a randomized clinical trial. *Trials*. 2024; 25(1): 536. doi: 10.1186/s13063-024-08389-y
82. Batalik L, Winnige P, Dosbaba F, et al. Home-based aerobic and resistance exercise interventions in cancer patients and survivors: A systematic review. *Cancers*. 2021; 13(8): 1915. doi: 10.3390/cancers13081915
83. Campbell KL, Winters-Stone KM, Wiskemann J, et al. Exercise guidelines for cancer survivors: consensus statement from international multidisciplinary roundtable. *Medicine & Science in Sports & Exercise*. 2019; 51(11): 2375-2390. doi: 10.1249/MSS.0000000000002116